

表 87 2016年導入患者, 年齢, 性別 (埼玉県)

| 年齢 | 男性 | 女性 | 合計 | 記載なし | 総計 |
|------|---------|---------|---------|------|---------|
| 5歳未満 | | | | | |
| (%) | | | | | |
| 5歳～ | 2 | 1 | 3 | | 3 |
| (%) | (0.1) | (0.2) | (0.1) | | (0.1) |
| 10歳～ | 1 | | 1 | | 1 |
| (%) | (0.1) | | (0.0) | | (0.0) |
| 15歳～ | 1 | | 1 | | 1 |
| (%) | (0.1) | | (0.0) | | (0.0) |
| 20歳～ | 2 | 2 | 4 | | 4 |
| (%) | (0.1) | (0.3) | (0.2) | | (0.2) |
| 25歳～ | 5 | 2 | 7 | | 7 |
| (%) | (0.3) | (0.3) | (0.3) | | (0.3) |
| 30歳～ | 12 | 3 | 15 | | 15 |
| (%) | (0.8) | (0.5) | (0.7) | | (0.7) |
| 35歳～ | 23 | 12 | 35 | | 35 |
| (%) | (1.5) | (1.9) | (1.6) | | (1.6) |
| 40歳～ | 56 | 22 | 78 | | 78 |
| (%) | (3.7) | (3.5) | (3.7) | | (3.7) |
| 45歳～ | 98 | 21 | 119 | | 119 |
| (%) | (6.5) | (3.4) | (5.6) | | (5.6) |
| 50歳～ | 91 | 26 | 117 | | 117 |
| (%) | (6.0) | (4.2) | (5.5) | | (5.5) |
| 55歳～ | 101 | 38 | 139 | | 139 |
| (%) | (6.7) | (6.1) | (6.5) | | (6.5) |
| 60歳～ | 125 | 40 | 165 | | 165 |
| (%) | (8.3) | (6.4) | (7.7) | | (7.7) |
| 65歳～ | 238 | 88 | 326 | | 326 |
| (%) | (15.7) | (14.2) | (15.3) | | (15.3) |
| 70歳～ | 225 | 94 | 319 | | 319 |
| (%) | (14.9) | (15.1) | (14.9) | | (14.9) |
| 75歳～ | 232 | 115 | 347 | | 347 |
| (%) | (15.3) | (18.5) | (16.3) | | (16.3) |
| 80歳～ | 207 | 91 | 298 | | 298 |
| (%) | (13.7) | (14.7) | (14.0) | | (14.0) |
| 85歳～ | 76 | 50 | 126 | | 126 |
| (%) | (5.0) | (8.1) | (5.9) | | (5.9) |
| 90歳～ | 19 | 13 | 32 | | 32 |
| (%) | (1.3) | (2.1) | (1.5) | | (1.5) |
| 95歳～ | | 3 | 3 | | 3 |
| (%) | | (0.5) | (0.1) | | (0.1) |
| 合計 | 1,514 | 621 | 2,135 | | 2,135 |
| (%) | (100.0) | (100.0) | (100.0) | | (100.0) |
| 不明 | | | | | |
| 記載なし | | | | | |
| 総計 | 1,514 | 621 | 2,135 | | 2,135 |
| 平均 | 67.45 | 69.83 | 68.14 | | 68.14 |
| 標準偏差 | 13.68 | 13.66 | 13.71 | | 13.71 |

数値下のかっこ内は列方向の合計に対する%です。

##A0022##

(C)Japanese Society for Dialysis Therapy

2016年調査