

表 127 2016年導入患者、年齢、原疾患別(女性)

原疾患	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差			
慢性系球体腎炎 (%)					1	3	8	20	40	64	78	97	124	174	296	291	336	344	225	68	10	2,179		2,179	70.41	13.94			
慢性腎盂腎炎 (%)						(12.5)	(17.6)	(22.9)	(23.8)	(29.9)	(25.7)	(18.8)	(19.5)	(19.8)	(17.7)	(18.2)	(19.3)	(17.9)	(17.3)	(18.0)	(18.0)	(18.2)	(18.6)	(18.6)	120	69.17	14.27		
急速進行性系球体腎炎 (%)					1	2	1	1	3	8	7	4	8	24	15	17	11	18	1							(1.0)			
妊娠腎/妊娠中毒症 (%)						(25.0)	(5.9)	(5.7)	(1.2)	(0.7)	(0.8)	(0.7)	(0.6)	(1.0)	(2.1)	(1.6)	(2.3)	(2.2)	(2.6)	(2.5)	(1.3)	(3.6)	(2.0)	(2.0)	233	73.96	12.66		
その他の分類不能の腎炎 (%)	1						(25.0)			(1.5)	(0.4)	(0.7)	(0.4)	(0.8)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)					24	60.33	13.97		
多発性囊胞腎 (%)	1						(25.0)			(11.8)	(2.4)	(0.8)	(1.0)	(0.2)	(0.5)	(0.6)	(0.3)	(0.8)	(0.5)	(0.4)	(0.4)	(0.3)	(1.8)	(0.5)	(0.5)	(0.5)	(0.5)	(0.5)	
腎硬化症 (%)										2	2	4	1	3	6	5	12	9	8	5	1	1	62	62	67.02	18.22			
悪性高血圧 (%)										3	5	8	29	41	58	56	74	49	43	32	18	6							
糖尿病性腎症 (%)										(3.6)	(3.7)	(3.2)	(7.0)	(8.2)	(9.3)	(5.7)	(4.6)	(3.2)	(2.3)	(1.6)	(1.4)	(1.6)		423	64.90	12.40			
SLE腎炎 (%)	1									(5.7)	(3.6)	(4.5)	(3.2)	(5.3)	(6.2)	(4.6)	(7.4)	(10.1)	(11.9)	(15.6)	(19.8)	(22.9)	(28.0)	(27.3)	(13.7)	(13.7)	(13.7)		
アミロイド腎 (%)										2	3	6	8	22	31	29	73	164	180	292	393	287	106	15	1,611	77.21	10.67		
痛風腎 (%)										1	2	5	6	7	4	3	11	10	14	13	8	3	1	89	68.01	16.14			
先天性代謝異常による腎不全 (%)										1	2	4	23	44	93	177	199	278	438	711	627	717	623	296	70	8	4,311	69.67	12.31
腎・尿路結核 (%)										1	2	4	(1.2)	(1.5)	(2.0)	(1.4)	(1.4)	(0.6)	(0.3)	(0.7)	(0.7)	(0.7)	(0.6)	(0.8)	(1.8)	(0.8)	(0.8)	(0.8)	
腎・尿路結石 (%)										1	2	4	(11.8)	(11.4)	(27.4)	(32.8)	(37.3)	(42.7)	(40.0)	(44.3)	(44.5)	(43.8)	(41.6)	(38.2)	(31.4)	(23.6)	(18.5)	(14.5)	(36.7)
閉塞性尿路障害 (%)										1	2	4	(5.9)	(5.7)	(3.6)	(3.0)	(3.2)	(3.6)	(1.8)	(1.9)	(1.3)	(0.9)	(1.1)	(0.9)	(0.7)	(0.2)	(0.3)	(1.1)	133
骨髄腫 (%)										1	2	4	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)	(2.5)
腎形成不全 (%)	1									1	2	3	(25.0)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)	(12.5)
不明 (%)										1	2	3	(75.0)	(75.0)	(50.0)	(12.5)	(35.3)	(8.6)	(10.7)	(7.5)	(5.6)	(4.1)	(4.8)	(3.3)	(4.4)	(3.6)	(4.8)	(5.1)	(5.3)
再導入 (%)										1	3	1	(12.5)	(25.0)	(5.9)	(25.7)	(15.5)	(6.7)	(11.2)	(10.6)	(11.5)	(9.1)	(11.0)	(11.4)	(11.9)	(13.9)	(17.8)	(21.2)	(25.5)
その他 (%)	3	3	2	1	6	3	9	10	14	17	24	21	43	55	54	91	101	70	20	4	551	551	70.08	17.35					
合計 (%)	4	4	4	8	17	35	84	134	249	415	497	627	984	1,625	1,506	1,877	1,987	1,252	378	55	11,744		11,744	71.19	13.47				
記載なし																													
総計	4	4	4	8	17	35	84	134	249	415	497	627	984	1,625	1,506	1,877	1,987	1,252	378	55	11,744		11,744	71.19	13.47				

数値下のかっこ内は列方向の合計に対する%です。

##A0062##

(C)Japanese Society for Dialysis Therapy

2016年調査