

表 128 2016年導入患者, 原疾患別, 平均年齢

原疾患	人数	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	6,186		6,186	69.65	13.78
(%)	(16.6)		(16.6)		
慢性腎盂腎炎	258		258	68.53	14.94
(%)	(0.7)		(0.7)		
急速進行性糸球体腎炎	475		475	73.30	12.18
(%)	(1.3)		(1.3)		
妊娠腎/妊娠中毒症	24		24	60.33	13.97
(%)	(0.1)		(0.1)		
その他の分類不能の腎炎	169		169	63.76	18.95
(%)	(0.5)		(0.5)		
多発性嚢胞腎	972		972	63.38	13.31
(%)	(2.6)		(2.6)		
腎硬化症	5,285		5,285	75.06	11.66
(%)	(14.2)		(14.2)		
悪性高血圧	290		290	65.91	16.53
(%)	(0.8)		(0.8)		
糖尿病性腎症	16,103		16,103	67.30	12.22
(%)	(43.2)		(43.2)		
SLE腎炎	219		219	63.48	14.71
(%)	(0.6)		(0.6)		
アミロイド腎	80		80	67.13	12.32
(%)	(0.2)		(0.2)		
痛風腎	72		72	67.47	11.48
(%)	(0.2)		(0.2)		
先天性代謝異常による腎不全	23		23	50.83	22.41
(%)	(0.1)		(0.1)		
腎・尿路結核	9		9	74.56	8.83
(%)	(0.0)		(0.0)		
腎・尿路結石	55		55	69.64	11.73
(%)	(0.1)		(0.1)		
腎・尿路腫瘍	191		191	72.23	10.27
(%)	(0.5)		(0.5)		
閉塞性尿路障害	90		90	70.58	12.37
(%)	(0.2)		(0.2)		
骨髄腫	153		153	71.35	10.98
(%)	(0.4)		(0.4)		
腎形成不全	47		47	40.17	23.94
(%)	(0.1)		(0.1)		
不明	4,759	2	4,761	72.26	13.41
(%)	(12.8)	(100.0)	(12.8)		
再導入	238		238	57.85	15.84
(%)	(0.6)		(0.6)		
その他	1,552		1,552	69.49	16.07
(%)	(4.2)		(4.2)		
合計	37,250	2	37,252	69.40	13.36
(%)	(100.0)	(100.0)	(100.0)		
記載なし					
総計	37,250	2	37,252	69.40	13.36

数値下のかっこ内は列方向の合計に対する%です。

##A0063##

(C)Japanese Society for Dialysis Therapy

2016年調査