

表 134 2016年導入患者、治療方法(新分類)、原疾患別

原疾患	施設血液透析	Off-line HDF	On-line HDF	Push/Pull HDF	ハイオフィードション	IHDF	血液濾過	血液吸着透析(リクセル等使用)	在宅血液透析	腹膜透析(CAPD)	腹膜透析(APD)	透析離脱	生体腎移植(親から)	生体腎移植(祖父母から)	生体腎移植(兄弟から)	生体腎移植(子から)	生体腎移植(それ以外の血縁者から)	生体腎移植(配偶者から)	生体腎移植(上記以外から)	臓器移植(生体腎が献腎か不明の場合)	献腎移植	合計	記載なし	総計	
慢性系球体腎炎 (%)	4,629 (79.4)	28 (0.5)	498 (8.5)			156 (2.7)		1 (0.0)	4 (0.1)	251 (4.3)	150 (2.6)	36 (0.6)	11 (0.2)	31 (0.5)		10 (0.2)		20 (0.3)	3 (0.1)			5,826 (100.0)	358 (100.0)	6,186	
慢性腎孟腎炎 (%)	197 (82.4)		21 (8.8)			4 (1.7)				9 (3.8)	4 (1.7)	2 (0.8)	1 (0.4)					1 (0.4)				239 (100.0)	19 (100.0)	258	
急速進行性系球体腎炎 (%)	352 (86.3)	2 (0.5)	25 (6.1)			9 (2.2)				7 (1.7)	4 (1.0)	9 (2.2)										406 (100.0)	67 (100.0)	475	
妊娠腎/妊娠中毒症	16		4																			24 (100.0)	24 (100.0)	24	
その他の分類不能の腎炎 (%)	128 (81.5)		10 (6.4)			2 (1.3)				4 (2.5)	6 (3.8)	5 (3.2)	1 (0.6)			1 (4.2)						157 (100.0)	12 (100.0)	169	
多発性囊胞腎 (%)	756 (81.3)	4 (0.4)	73 (7.8)			21 (2.3)				35 (3.8)	19 (2.0)	6 (0.6)		5 (0.5)		2 (0.2)	1 (0.1)	7 (0.8)	1 (0.1)	1 (0.1)		930 (100.0)	42 (100.0)	972	
腎硬化症 (%)	4,049 (81.8)	25 (0.5)	443 (8.9)	1 (0.0)	1 (0.0)	146 (2.9)				1 (0.0)	174 (3.5)	76 (1.5)	7 (0.1)	7 (0.1)		3 (0.1)		11 (0.2)				4,951 (100.0)	334 (100.0)	5,285	
悪性高血圧 (%)	223 (80.2)	1 (0.4)	20 (7.2)			13 (4.7)					14 (5.0)	4 (1.4)	1 (0.4)				1 (0.1)					278 (100.0)	12 (100.0)	290	
糖尿病性腎症 (%)	12,608 (82.6)	67 (0.4)	1,388 (9.1)	1 (0.0)	1 (0.0)	449 (2.9)	1 (0.0)			1 (0.0)	413 (2.7)	223 (1.5)	52 (0.3)	28 (0.2)	6 (0.0)		2 (0.0)	2 (0.0)	21 (0.1)	2 (0.0)	3 (0.0)	15,268 (100.0)	835 (100.0)	16,103	
SLE腎炎	160		19			5 (2.5)					6 (3.0)	5 (2.5)	2 (1.0)	1 (0.5)			1 (0.5)					199 (100.0)	20 (100.0)	219	
アミロイド腎 (%)	49 (70.0)	3 (4.3)	7 (10.0)			3 (4.3)					5 (7.1)	3 (4.3)										70 (100.0)	10 (100.0)	80	
痛風腎 (%)	57 (81.4)		8 (11.4)								3 (4.3)			1 (1.4)				1 (1.4)				70 (100.0)	2 (100.0)	72	
先天性代謝異常による腎不全 (%)	19 (82.6)		2 (8.7)			1 (4.3)					1 (4.3)											23 (100.0)	23 (100.0)	23	
腎・尿路結核 (%)	6 (75.0)		1 (12.5)								1 (12.5)											8 (100.0)	1 (100.0)	9	
腎・尿路結石 (%)	43 (81.1)		3 (5.7)			2 (3.8)					4 (7.5)				1 (1.9)							53 (100.0)	2 (100.0)	55	
腎・尿路腫瘍 (%)	147 (85.0)	1 (0.6)	18 (10.4)			5 (2.9)					1 (0.6)	1 (0.6)										173 (100.0)	18 (100.0)	191	
閉塞性尿路障害 (%)	71 (85.5)		1 (4.8)			2 (2.4)					2 (2.4)	1 (1.2)		1 (1.2)	1 (1.2)							83 (100.0)	7 (100.0)	90	
骨髄腫 (%)	116 (87.9)	2 (1.5)	3 (2.3)			2 (1.5)					1 (0.8)	3 (2.3)		2 (3.8)	5 (3.0)							132 (100.0)	21 (100.0)	153	
腎形成不全 (%)	25 (54.3)		6 (13.0)									9 (19.6)	2 (4.3)		3 (6.5)							46 (100.0)	1 (100.0)	47	
不明 (%)	3,522 (82.7)	19 (0.4)	345 (8.1)	1 (0.0)	118 (2.8)					1 (0.0)	129 (3.0)	61 (1.4)	30 (0.7)	12 (0.3)	11 (0.3)		1 (0.0)	8 (0.2)	2 (0.0)	1 (0.0)		4,260 (100.0)	501 (100.0)	4,761	
再導入 (%)	159 (71.9)	1 (0.5)	22 (10.0)			7 (3.2)					2 (0.9)	13 (5.9)	2 (0.9)	2 (0.9)	5 (2.3)		2 (0.9)	5 (2.3)	1 (0.5)			221 (100.0)	17 (100.0)	238	
その他 (%)	1,122 (82.4)	14 (1.0)	103 (7.6)			31 (2.3)					1 (0.1)	37 (2.7)	25 (1.8)	2 (0.1)	18 (1.3)	7 (0.5)			1 (0.1)				1,361 (100.0)	191 (100.0)	1,552
合計	28,454	168 (81.8)	3,023 (0.5)	2 (8.7)	3 (0.0)	976 (2.8)	1 (0.0)	1 (0.0)	10 (0.0)	1,111 (3.2)	598 (1.7)	147 (0.4)	94 (0.3)	79 (0.2)		21 (0.1)	4 (0.0)	1 (0.0)	75 (0.2)	6 (0.0)	8 (0.0)		34,782 (100.0)	2,470 (100.0)	37,252
記載なし (%)																									
総計	28,454	168 (81.8)	3,023 (0.5)	2 (8.7)	3 (0.0)	976 (2.8)	1 (0.0)	1 (0.0)	10 (0.0)	1,111 (3.2)	598 (1.7)	147 (0.4)	94 (0.3)	79 (0.2)		21 (0.1)	4 (0.0)	1 (0.0)	75 (0.2)	6 (0.0)	8 (0.0)		34,782 (100.0)	2,470 (100.0)	37,252

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

#A0069#

(C)Japanese Society for Dialysis Therapy

2016年調査