

表 137 2016年導入患者死亡原因分類(年齢との関係)

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差	
心不全										2	5	4	14	13	38	57	82	131	107	39	6	498		498	79.01	9.52	
(%)										(15.4)	(27.8)	(9.3)	(21.2)	(12.4)	(17.8)	(21.9)	(20.7)	(23.6)	(25.9)	(28.7)	(23.1)	(22.1)		(22.1)			
脳血管障害										5	2	3	11	7	15	16	21	26	21	3	3	133		133	73.84	12.46	
(%)										(38.5)	(11.1)	(7.0)	(16.7)	(6.7)	(7.0)	(6.2)	(5.3)	(4.7)	(5.1)	(2.2)	(11.5)	(5.9)		(5.9)			
感染症										2	5	2	9	12	30	41	67	114	156	115	36	6	595		595	78.06	9.79
(%)										(50.0)	(38.5)	(11.1)	(20.9)	(18.2)	(28.6)	(19.2)	(25.8)	(28.7)	(28.1)	(27.8)	(26.5)	(23.1)	(26.4)	(26.4)			
消化管出血										1	1	4	5	6	4	17	6	4	4	4	48		48	77.13	9.65		
(%)										(2.3)	(1.5)	(3.8)	(2.3)	(2.3)	(1.0)	(3.1)	(1.5)	(2.9)	(2.9)	(2.1)		(2.1)					
悪性腫瘍										2	8	8	20	51	40	49	53	34	9		274		274	74.54	9.58		
(%)										(11.1)	(18.6)	(12.1)	(19.0)	(23.8)	(15.4)	(12.3)	(9.5)	(8.2)	(6.6)		(12.2)		(12.2)				
悪液質/尿毒症										1	2	2	5	10	13	30	23	12	98		98		80.63	9.18			
(%)										(7.7)	(4.7)	(1.9)	(2.3)	(3.8)	(3.3)	(5.4)	(5.6)	(8.8)	(4.4)		(4.4)						
心筋梗塞										1	1	2	6	7	13	17	12	3	4	66		66		79.23	9.62		
(%)										(5.6)	(1.5)	(1.5)	(1.9)	(2.8)	(2.7)	(3.3)	(3.1)	(2.9)	(2.2)	(15.4)	(2.9)	(2.9)					
カリウム中毒/頻死										1	4	1	5	3	5	6	3			28		28		72.75	10.41		
(%)										(2.3)	(6.1)	(1.0)	(2.3)	(1.2)	(1.3)	(1.1)	(0.7)			(1.2)		(1.2)					
肝硬変症										2	4	1	4	5	8	9	5	1		39		39		69.74	10.71		
(%)										(11.1)	(9.3)	(1.5)	(3.8)	(2.3)	(3.1)	(2.3)	(0.9)	(0.2)		(1.7)		(1.7)					
自殺/拒否										1	1	2	1	5	8	2			20		20		76.45	7.70			
(%)										(1.5)	(1.0)	(0.9)	(0.4)	(1.3)	(1.4)	(0.5)			(0.9)		(0.9)						
腸閉塞										1	1	2	4	11	5	1	1	1	26		26		81.38	7.09			
(%)										(1.0)	(0.5)	(0.8)	(1.0)	(2.0)	(1.2)	(0.7)	(3.8)	(1.2)		(1.2)		(1.2)					
肺血栓/肺塞栓										1	3					1			1	6		6		70.83	15.09		
(%)										(5.6)	(1.5)	(1.5)	(2.9)			(0.2)			(3.8)	(0.3)		(0.3)					
災害・事故死										1	1	1								3		3		57.00	8.00		
(%)										(5.6)	(1.5)	(1.5)	(0.5)							(0.1)		(0.1)					
その他										7	5	11	21	25	46	57	52	20	2	246		246		78.56	9.65		
(%)										(16.3)	(7.6)	(10.5)	(9.8)	(9.6)	(11.6)	(10.3)	(12.6)	(14.7)	(7.7)	(10.9)		(10.9)					
不明	1									2	3	4	6	6	18	18	32	38	32	9	3	172		172		76.34	12.46
(%)	(100.0)									(50.0)	(16.7)	(9.3)	(9.1)	(5.7)	(8.4)	(6.9)	(8.1)	(6.8)	(7.7)	(6.6)	(11.5)	(7.6)	(7.6)				
合計	1									4	13	18	43	66	105	214	260	397	556	413	136	26	2,252		2,252	77.41	10.29
(%)	(100.0)									(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)				
記載なし										4	13	18	43	66	105	214	260	397	556	413	136	26	2,252		2,252	77.41	10.29
総計	1																										

数値下のかっこ内は列方向の合計に対する%です。

##A072##

(C)Japanese Society for Dialysis Therapy

2016年調査