

表 138 2016年導入患者死亡原因分類(透析期間との関係)

死因	3ヶ月未満	3ヶ月～	6ヶ月～	合計	不明	記載なし	総計
心不全	270	137	90	497	1		498
(%)	(22.6)	(21.8)	(21.1)	(22.1)	(33.3)		(22.1)
脳血管障害	71	38	24	133			133
(%)	(6.0)	(6.0)	(5.6)	(5.9)			(5.9)
感染症	344	153	98	595			595
(%)	(28.8)	(24.3)	(23.0)	(26.5)			(26.4)
消化管出血	28	13	7	48			48
(%)	(2.3)	(2.1)	(1.6)	(2.1)			(2.1)
悪性腫瘍	121	84	69	274			274
(%)	(10.1)	(13.4)	(16.2)	(12.2)			(12.2)
悪液質/尿毒症	45	34	19	98			98
(%)	(3.8)	(5.4)	(4.4)	(4.4)			(4.4)
心筋梗塞	44	17	5	66			66
(%)	(3.7)	(2.7)	(1.2)	(2.9)			(2.9)
カリウム中毒/頓死	9	9	10	28			28
(%)	(0.8)	(1.4)	(2.3)	(1.2)			(1.2)
肝硬変症	23	10	6	39			39
(%)	(1.9)	(1.6)	(1.4)	(1.7)			(1.7)
自殺/拒否	9	5	6	20			20
(%)	(0.8)	(0.8)	(1.4)	(0.9)			(0.9)
腸閉塞	13	11	2	26			26
(%)	(1.1)	(1.7)	(0.5)	(1.2)			(1.2)
肺血栓/肺塞栓	3	2	1	6			6
(%)	(0.3)	(0.3)	(0.2)	(0.3)			(0.3)
災害・事故死	1	1	1	3			3
(%)	(0.1)	(0.2)	(0.2)	(0.1)			(0.1)
その他	137	69	40	246			246
(%)	(11.5)	(11.0)	(9.4)	(10.9)			(10.9)
不明	75	46	49	170	2		172
(%)	(6.3)	(7.3)	(11.5)	(7.6)	(66.7)		(7.6)
合計	1,193	629	427	2,249	3		2,252
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし							
総計	1,193	629	427	2,249	3		2,252

数値下のカッコ内は列方向の合計に対する%です。

##A0073##

(C)Japanese Society for Dialysis Therapy

2016年調査