

表 141 2016年導入患者死亡原因分類(年齢別,原疾患別)原疾患=慢性腎盂腎炎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)															1 (100.0)			3 (75.0)			4 (22.2)			4 (22.2)	83.75	7.27
脳血管障害 (%)																										
感染症 (%)											1 (33.3)					3 (75.0)			1 (100.0)		5 (27.8)			5 (27.8)	76.00	14.76
消化管出血 (%)																										
悪性腫瘍 (%)											2 (66.7)		2 (100.0)			1 (25.0)	1 (50.0)				6 (33.3)			6 (33.3)	64.83	12.35
悪液質/尿毒症 (%)																		1 (25.0)			1 (5.6)			1 (5.6)	89.00	
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腫瘍塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)														1 (100.0)							1 (5.6)			1 (5.6)	67.00	
不明 (%)																	1 (50.0)				1 (5.6)			1 (5.6)	80.00	
合計 (%)											3 (100.0)		2 (100.0)	1 (100.0)	1 (100.0)	4 (100.0)	2 (100.0)	4 (100.0)	1 (100.0)		18 (100.0)			18 (100.0)	74.44	13.29
記載なし																										
脚計											3		2	1	1	4	2	4	1		18			18	74.44	13.29

数値下のかっこ内は列方向の合計に対する%です。  
##A0076##  
(C)Japanese Society for Dialysis Therapy  
2016年調査