

表 144 2016年導入患者死亡原因分類(年齢別,原疾患別)原疾患 = その他分類不能の腎炎

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)																1 (33.3)					1 (7.1)			1 (7.1)	77.00	
脳血管障害 (%)															1 (33.3)						1 (7.1)			1 (7.1)	74.00	
感染症 (%)													1 (100.0)			1 (33.3)	2 (66.7)	1 (33.3)			5 (35.7)			5 (35.7)	77.60	8.91
消化管出血 (%)																										
悪性腫瘍 (%)																2 (66.7)	1 (33.3)				3 (21.4)			3 (21.4)	73.67	2.52
悪液質/尿毒症 (%)																			1 (33.3)		1 (7.1)			1 (7.1)	89.00	
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腫瘍塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																	1 (33.3)		1 (33.3)					2 (14.3)	84.00	2.83
不明 (%)																				1 (100.0)	1 (7.1)			1 (7.1)	95.00	
合計 (%)													1 (100.0)		3 (100.0)	3 (100.0)	3 (100.0)	3 (100.0)		1 (100.0)	14 (100.0)			14 (100.0)	79.43	8.16
記載なし																										
脚計													1		3	3	3	3		1	14			14	79.43	8.16

数値下のかっこ内は列方向の合計に対する%です。
##A0079##
(C)Japanese Society for Dialysis Therapy
2016年調査