

表 145 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患 = 多発性囊胞腎

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)																					3	3	70.33	22.94			
脳血管障害 (%)																					(10.7)	3	78.67	11.59			
感染症 (%)																					(10.7)	7	77.57	8.06			
消化管出血 (%)																					(25.0)	1	81.00				
悪性腫瘍 (%)																					(3.6)	6	68.67	11.24			
悪液質/尿毒症 (%)																					(21.4)	3	82.00	10.58			
カリウム中毒/頻死 (%)																					(10.7)						
心筋梗塞 (%)																					1	1	81.00				
肝硬変症 (%)																					(3.6)	1	81.00				
自殺/拒否 (%)																					(3.6)	1	81.00				
腸閉塞 (%)																					1	1	81.00				
肺血栓/肺塞栓 (%)																					(3.6)	1	81.00				
災害・事故死 (%)																					2	2	66.50	9.19			
その他 (%)																					(7.1)	2	84.00	7.07			
不明 (%)																					(7.1)	2	84.00	7.07			
合計 (%)									1 (100.0)	1 (100.0)	2 (100.0)	3 (100.0)	5 (100.0)	4 (100.0)	5 (100.0)	4 (100.0)	5 (100.0)	4 (100.0)	3 (100.0)	28 (100.0)			28 (100.0)	75.39	11.57		
記載なし																											
総計									1	1	2	3	5	4	5	4	5	4	3	28			28	75.39	11.57		

数値下のかっこ内は列方向の合計に対する%です。

##A080##

(C)Japanese Society for Dialysis Therapy

2016年調査