

表 147 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患 = 惠性高血圧

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)															1 (50.0)		1 (33.3)		1 (100.0)	1 (27.3)	3	3 (27.3)	84.67	13.05		
脳血管障害 (%)															1 (50.0)		1 (33.3)		1 (100.0)	1 (9.1)	1	1 (9.1)	70.00			
感染症 (%)																										
消化管出血 (%)																										
悪性腫瘍 (%)																										
悪液質/尿毒症 (%)																										
カリウム中毒/頻死 (%)																										
心筋梗塞 (%)																										
カリウム中毒/頻死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																										
不明 (%)																										
合計 (%)															2 (100.0)	2 (100.0)	2 (100.0)	3 (100.0)	1 (100.0)	1 (100.0)	11 (100.0)		11 (100.0)	82.09	8.49	
記載なし																										
総計															2	2	2	3	1	1	11		11	82.09	8.49	

数値下のかっこ内は列方向の合計に対する%です。

##A082##
(C)Japanese Society for Dialysis Therapy
2016年調査