

表 150 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患 = アミロイド腎

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)										1 (50.0)					1 (50.0)	1 (33.3)	2 (33.3)				5 (27.8)			5 (27.8)	72.60	14.28
脳血管障害 (%)																								5 (27.8)	67.20	16.02
感染症 (%)																								2 (11.1)	80.50	0.71
消化管出血 (%)																										
悪性腫瘍 (%)																										
悪液質/尿毒症 (%)																										
心筋梗塞 (%)																										
カリウム中毒/頻死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																										
不明 (%)																										
合計 (%)									1 (100.0)	2 (100.0)		1 (100.0)	1 (100.0)	2 (100.0)	2 (100.0)	3 (100.0)	6 (100.0)				18 (100.0)			18 (100.0)	70.11	13.56
記載なし																										
総計									1	2		1	1	2	2	3	6				18			18	70.11	13.56

数値下のかっこ内は列方向の合計に対する%です。

##A085##
(C)Japanese Society for Dialysis Therapy
2016年調査