

表 154 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患 = 腎・尿路結石

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)																					1	(25.0)		1	80.00	
脳血管障害 (%)																					1	(25.0)		1	62.00	
感染症 (%)																					1	(25.0)		1	79.00	1.41
消化管出血 (%)																					2	(50.0)		2	(50.0)	
悪性腫瘍 (%)																					1	(100.0)		1	(100.0)	
悪液質/尿毒症 (%)																					1	(100.0)		1	(100.0)	
心筋梗塞 (%)																					1	(100.0)		1	(100.0)	
カリウム中毒/頓死 (%)																					1	(100.0)		1	(100.0)	
肝硬変症 (%)																					1	(100.0)		1	(100.0)	
自殺/拒否 (%)																					1	(100.0)		1	(100.0)	
腸閉塞 (%)																					1	(100.0)		1	(100.0)	
肺血栓/肺塞栓 (%)																					1	(100.0)		1	(100.0)	
災害・事故死 (%)																					1	(100.0)		1	(100.0)	
その他 (%)																					1	(100.0)		1	(100.0)	
不明 (%)																					1	(100.0)		1	(100.0)	
合計 (%)																					1	(100.0)		1	(100.0)	
記載なし																					1	(100.0)		1	(100.0)	
総計																					1	(100.0)		1	(100.0)	

数値下のかっこ内は列方向の合計に対する%です。

##A0089##
(C)Japanese Society for Dialysis Therapy
2016年調査