

表 155 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患=腎・尿路腫瘍

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差	
心不全 (%)																											
脳血管障害 (%)																											
感染症 (%)																											
消化管出血 (%)																											
悪性腫瘍 (%)																											
悪液質/尿毒症 (%)																											
カリウム中毒/頓死 (%)																											
肝硬変症 (%)																											
自殺/拒否 (%)																											
腸閉塞 (%)																											
肺血栓/肺塞栓 (%)																											
災害・事故死 (%)																											
その他 (%)																											
不明 (%)																											
合計 (%)											1 (100.0)	1 (100.0)	2 (100.0)	1 (100.0)	3 (100.0)	2 (100.0)	4 (100.0)	2 (100.0)	16 (100.0)	16 (100.0)				16 (100.0)	77.31	11.03	
記載なし											1	1	2	1	3	2	4	2	16								
総計																									16	77.31	11.03

数値下の括弧内は列方向の合計に対する%です。

#A0090##

(C)Japanese Society for Dialysis Therapy

2016年調査