

表 156 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患 = 閉塞性尿路障害

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)																					2		2	77.00	1.41	
脳血管障害 (%)																					(15.4)		(15.4)			
感染症 (%)																					8		8	74.88	10.44	
消化管出血 (%)																					1		1	79.00		
悪性腫瘍 (%)																					1		1	7.7		
悪液質/尿毒症 (%)																					1		1	68.00		
心筋梗塞 (%)																					1		1	7.7		
カリウム中毒/頻死 (%)																					1		1	7.7		
肝硬変症 (%)																					1		1	7.7		
自殺/拒否 (%)																					1		1	7.7		
腸閉塞 (%)																					1		1	7.7		
肺血栓/肺塞栓 (%)																					1		1	7.7		
災害・事故死 (%)																					1		1	7.7		
その他 (%)																					1		1	7.7		
不明 (%)																					1		1	7.7		
合計 (%)																					13		13	74.92	8.36	
記載なし																										
総計																					13		13	74.92	8.36	

数値下のかっこ内は列方向の合計に対する%です。

##A0091##

(C)Japanese Society for Dialysis Therapy

2016年調査