

表 157 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患=骨髄腫

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差		
心不全 (%)											1 (100.0)		1 (25.0)	1 (10.0)							2 (6.3)			2 (6.3)	66.00	2.83		
脳血管障害 (%)												1 (100.0)									1 (3.1)			1 (3.1)	44.00			
感染症 (%)													1 (100.0)	1 (10.0)								4 (12.5)			4 (12.5)	72.50	14.98	
消化管出血 (%)													1 (100.0)	3 (75.0)	7 (70.0)	2 (66.7)	2 (100.0)	1 (60.0)	3 (60.0)	3 (60.0)			21 (65.6)			21 (65.6)	72.00	10.34
悪性腫瘍 (%)														1 (33.3)							2 (6.3)			2 (6.3)	79.50	13.44		
悪液質/尿毒症 (%)																												
心筋梗塞 (%)																												
カリウム中毒/鎮死 (%)																												
肝硬変症 (%)																												
自殺/拒否 (%)																												
腸閉塞 (%)																												
肺血栓/肺塞栓 (%)																												
災害・事故死 (%)																												
その他 (%)																					1 (3.1)			1 (3.1)	66.00			
不明 (%)																												
合計 (%)											1 (100.0)	1 (100.0)	1 (100.0)	4 (100.0)	10 (100.0)	3 (100.0)	2 (100.0)	5 (100.0)	5 (100.0)		32 (100.0)			32 (100.0)	71.44	11.52		
記載なし											1	1	1	4	10	3	2	5	5									
総計																					32			32	71.44	11.52		

数値下のかっこ内は列方向の合計に対する%です。

##A0092##

(C)Japanese Society for Dialysis Therapy

2016年調査