

表 160 2016年導入患者死亡原因分類(年齢別、原疾患別)原疾患 = 移植後再導入

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)											(50.0)	(50.0)	1 (33.3)	1 (33.3)	1 (50.0)	2 (50.0)					5 (27.8)			5 (27.8)	70.20	16.86
脳血管障害 (%)											1 (100.0)	1 (50.0)	1 (50.0)	1 (33.3)	2 (50.0)					2 (11.1)			2 (11.1)	57.50	7.78	
感染症 (%)																				5 (27.8)			5 (27.8)	61.80	12.28	
消化管出血 (%)																				1 (5.6)			1 (5.6)	64.00		
悪性腫瘍 (%)																				1 (5.6)			1 (5.6)	72.00		
悪液質/尿毒症 (%)																										
心筋梗塞 (%)																										
カリウム中毒/頓死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																										
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)																					3 (16.7)			3 (16.7)	73.67	5.03
不明 (%)																				1 (5.6)			1 (5.6)	79.00		
合計 (%)									1 (100.0)	2 (100.0)	2 (100.0)	3 (100.0)	2 (100.0)	4 (100.0)	2 (100.0)					18 (100.0)			18 (100.0)	67.28	12.19	
記載なし																										
総計									1	2	2	3	2	4	2					18			18	67.28	12.19	

数値下のかっこ内は列方向の合計に対する%です。

##A0095##  
(C)Japanese Society for Dialysis Therapy  
2016年調査