

表 162 2016年導入患者死亡原因分類(治療方法別)(2016年末)

死因	施設血液透析	血液透析濾過	血液濾過	血液吸着透析	在宅血液透析	腹膜透析	離脱	生体腎移植	献腎移植	献腎移植 (生体腎か 献腎か不明 の場合)	合計	記載なし	総計
心不全	197	15				5					217	281	498
(%)	(22.5)	(31.9)				(31.3)					(23.1)	(21.4)	(22.1)
脳血管障害	52	3				1					56	77	133
(%)	(5.9)	(6.4)				(6.3)					(6.0)	(5.9)	(5.9)
感染症	240	8				2					250	345	595
(%)	(27.4)	(17.0)				(12.5)					(26.7)	(26.3)	(26.4)
消化管出血	19	1									20	28	48
(%)	(2.2)	(2.1)									(2.1)	(2.1)	(2.1)
悪性腫瘍	110	5				1					116	158	274
(%)	(12.6)	(10.6)				(6.3)					(12.4)	(12.0)	(12.2)
悪液質/尿毒症	31					2					33	65	98
(%)	(3.5)					(12.5)					(3.5)	(4.9)	(4.4)
心筋梗塞	28										28	38	66
(%)	(3.2)										(3.0)	(2.9)	(2.9)
カリウム中毒/頓死	8	1									9	19	28
(%)	(0.9)	(2.1)									(1.0)	(1.4)	(1.2)
肝硬変症	9	1									10	29	39
(%)	(1.0)	(2.1)									(1.1)	(2.2)	(1.7)
自殺/拒否	9										9	11	20
(%)	(1.0)										(1.0)	(0.8)	(0.9)
腸閉塞	9					1					10	16	26
(%)	(1.0)					(6.3)					(1.1)	(1.2)	(1.2)
肺血栓/肺塞栓	5										5	1	6
(%)	(0.6)										(0.5)	(0.1)	(0.3)
災害・事故死	2										2	1	3
(%)	(0.2)										(0.2)	(0.1)	(0.1)
その他	91	2				1					94	152	246
(%)	(10.4)	(4.3)				(6.3)					(10.0)	(11.6)	(10.9)
不明	65	11				3					79	93	172
(%)	(7.4)	(23.4)				(18.8)					(8.4)	(7.1)	(7.6)
合計	875	47				16					938	1,314	2,252
(%)	(100.0)	(100.0)				(100.0)					(100.0)	(100.0)	(100.0)
記載なし													
総計	875	47				16					938	1,314	2,252

数値下のかっこ内は列方向の合計に対する%です。

##A0097##

(C)Japanese Society for Dialysis Therapy

2016年調査