

表 218 2016年末患者、年齢、原疾患別人数(女性)

原疾患	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差
慢性系球体腎炎	2	1	6	21	52	157	356	910	1,439	2,016	2,745	4,438	6,821	5,529	5,116	3,798	2,280	686	107	36,474			36,474	68.43	12.22	
(%)	(15.4)	(9.1)	(14.6)	(26.9)	(27.4)	(29.8)	(32.7)	(37.0)	(35.2)	(36.6)	(35.7)	(36.6)	(34.4)	(33.3)	(30.5)	(26.9)	(26.2)	(24.0)	(23.3)	(32.2)			(32.2)			
慢性腎盂腎炎			2	8	14	21	43	87	109	104	168	268	236	189	167	99	25	5	1,545			1,545	67.31	13.29		
(%)			(2.6)	(4.2)	(2.7)	(1.9)	(1.7)	(2.1)	(2.0)	(1.4)	(1.4)	(1.4)	(1.4)	(1.1)	(1.2)	(1.1)	(0.9)	(1.2)	(1.4)	(1.4)			(1.4)			
急速進行性系球体腎炎	2	1	5	6	7	12	24	41	37	72	115	200	182	205	187	111	29	1	1,237			1,237	70.47	12.97		
(%)	(18.2)	(2.4)	(6.4)	(3.2)	(1.3)	(1.1)	(1.0)	(1.0)	(0.7)	(0.9)	(0.9)	(1.0)	(1.1)	(1.2)	(1.3)	(1.0)	(0.2)	(1.1)	(1.1)	(1.1)			(1.1)			
妊娠腎/妊娠中毒症					3	9	28	78	100	157	229	353	231	168	98	25	4	1,483			1,483	65.76	10.27			
(%)					(0.6)	(0.8)	(1.1)	(1.9)	(1.8)	(2.0)	(1.9)	(1.8)	(1.4)	(1.0)	(0.7)	(0.3)	(0.1)	(1.3)	(1.3)			(1.3)				
その他の分類不能の腎炎	2		2	6	10	14	11	27	40	40	50	64	71	66	70	66	37	7	2	585			585	64.23	16.33	
(%)	(9.1)		(4.9)	(7.7)	(5.3)	(2.7)	(1.0)	(1.1)	(1.0)	(0.7)	(0.7)	(0.5)	(0.4)	(0.4)	(0.4)	(0.5)	(0.4)	(0.2)	(0.5)	(0.5)			(0.5)			
多発性囊胞腎	2	1	2	2	18	29	75	197	368	507	785	1,123	808	685	391	163	39	5	5,201			5,201	66.84	10.83		
(%)	(9.1)	(7.7)	(2.4)	(2.6)	(1.1)	(3.4)	(2.7)	(3.0)	(4.8)	(6.7)	(6.6)	(5.5)	(4.9)	(4.1)	(2.8)	(1.9)	(1.4)	(1.2)	(4.6)	(4.6)			(4.6)			
腎硬化症			1	4	9	22	29	77	156	213	289	551	1,206	1,263	1,854	2,206	1,818	719	148	10,565			10,565	76.54	11.18	
(%)			(2.4)	(5.1)	(4.7)	(4.2)	(2.7)	(3.1)	(3.8)	(3.9)	(3.8)	(3.8)	(4.5)	(6.1)	(7.6)	(11.0)	(15.6)	(20.9)	(25.2)	(34.2)	(34.2)	(9.3)	(9.3)			
悪性高血圧			1		3	5	18	28	47	57	67	84	122	100	116	113	63	36	9	873			873	68.63	14.39	
(%)			(2.4)		(1.6)	(0.9)	(1.7)	(1.1)	(1.2)	(1.0)	(0.9)	(0.7)	(0.6)	(0.6)	(0.7)	(0.8)	(0.7)	(1.3)	(2.1)	(0.8)	(0.8)	(0.8)	(0.8)			
糖尿病性腎症	1	1	1	16	104	280	679	1,163	1,639	2,490	3,858	6,854	5,835	5,737	4,529	2,320	654	55	36,216			36,216	69.73	11.52		
(%)	(9.1)	(2.4)	(1.3)	(8.4)	(19.7)	(25.7)	(27.6)	(28.5)	(29.8)	(32.4)	(31.8)	(34.6)	(35.1)	(34.1)	(32.1)	(26.7)	(22.9)	(22.7)	(32.0)	(32.0)			(32.0)			
SLE腎炎	1	2	10	18	39	93	146	142	185	226	238	174	113	80	32	9	1	1,509			1,509	61.31	12.98			
(%)	(2.4)	(2.6)	(5.3)	(3.4)	(3.6)	(3.8)	(3.6)	(2.6)	(2.4)	(1.9)	(1.2)	(1.0)	(0.7)	(0.6)	(0.4)	(0.3)	(0.2)	(1.3)	(1.3)			(1.3)				
アミロイド腎						1	2	5	14	16	29	61	58	29	28	11	2	1	257			257	69.56	9.78		
(%)						(0.1)	(0.1)	(0.1)	(0.3)	(0.2)	(0.2)	(0.3)	(0.3)	(0.2)	(0.2)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)			(0.2)			
痛風腎						1	1	1	11	10	7	17	24	15	22	13	10	1	133			133	67.72	12.62		
(%)						(0.2)	(0.1)	(0.0)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.0)	(0.1)	(0.1)	(0.1)			(0.1)			
先天性代謝異常による腎不全		4	2	6	10	14	9	6	9	17	9	4	4	4	2	1	111			111	53.89	16.31				
(%)		(5.1)	(1.1)	(1.1)	(0.9)	(0.6)	(0.3)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)	(0.1)			(0.1)				
腎・尿路結核																			83			83	73.83	8.79		
(%)																			(0.1)			(0.1)				
腎・尿路結石						1	1	6	7	11	21	40	27	27	28	21	5	2	197			197	71.66	11.63		
(%)						(0.2)	(0.1)	(0.2)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)			(0.2)			
腎・尿路腫瘍		1				3	7	13	12	14	19	14	25	44	19	38	21	17	6	217			217	72.69	11.54	
(%)		(2.4)				(1.6)	(1.3)	(1.2)	(0.5)	(0.3)	(0.3)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)			(0.1)			
閉塞性尿路障害	1																		253			253	64.32	16.24		
(%)		(7.7)																	(0.2)			(0.2)				
骨髄腫																			131			131	70.21	10.51		
(%)																			(0.1)			(0.1)				
腎形成不全	6	2	4	7	6	9	16	30	37	26	19	11	14	13	15	20	8	5	2	250			250	48.87	20.06	
(%)	(27.3)	(15.4)	(36.4)	(17.1)	(7.7)	(4.7)	(3.0)	(2.8)	(1.5)	(0.6)	(0.3)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)			(0.2)			
不明	1					7	8	20	61	105	200	379	435	627	1,043	1,750	1,536	1,880	1,893	1,347	525	84	11,901	72.07	12.86	
(%)	(4.5)					(17.1)	(10.3)	(10.5)	(11.6)	(9.7)	(8.1)	(9.3)	(7.9)	(8.2)	(8.6)	(8.8)	(9.2)	(11.2)	(13.4)	(15.5)	(18.4)	(19.4)	(10.5)			
再導入			3	6	10	17	37	69	73	91	104	116	116	30	32	21	16	5	746			746	57.06	13.61		
(%)			(7.3)	(7.7)	(5.3)	(3.2)	(3.4)	(2.8)	(1.8)	(1.7)	(1.4)	(1.0)	(0.6)	(0.2)	(0.1)	(0.2)	(0.2)	(0.2)	(0.7)	(0.7)			(0.7)			
その他	11	7	3	9	11	30	56	84	129	161	169	196	292	442	398	422	406	283	90	19	3,218			3,218	67.19	16.23
(%)	(50.0)	(53.8)	(27.3)	(22.0)	(14.1)	(15.8)	(10.6)	(7.7)	(5.2)	(3.9)	(3.1)	(2.6)	(2.4)	(2.2)	(2.4)	(2.5)	(2.9)	(3.3)	(3.2)	(4.4)	(2.8)			(2.8)		
合計	22	13	11	41	78	190	527	1,088	2,460	4,084	5,507	7,676	12,122	19,836	16,609	16,806	14,125	8,698	2,857	433	113,185			113,185	69.61	12.54
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)			(100.0)			
記載なし																										
総計	22	13	11	41	78	190	527	1,088	2,460	4,084	5,507	7,676	12,122	19,836	16,609	16,806	14,125	8,698	2,857	433	113,185			113,185	69.61	12.54

数値下の括弧内は列方向の合計に対する%です。

##A0153##

(C)Japanese Society for Dialysis Therapy

2016年調査