

表 219 2016年末患者，透析歴，原疾患別人数

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	12,060	15,697	19,324	14,760	11,606	8,192	4,885	2,950	1,641	628	91,743	40		91,783	11.41	9.54
(%)	(17.1)	(19.6)	(24.4)	(36.4)	(52.5)	(65.6)	(70.8)	(74.0)	(76.3)	(79.2)	(28.8)	(13.6)		(28.8)		
慢性腎盂腎炎	457	563	611	464	309	203	128	102	50	7	2,894	2		2,896	10.35	9.33
(%)	(0.6)	(0.7)	(0.8)	(1.1)	(1.4)	(1.6)	(1.9)	(2.6)	(2.3)	(0.9)	(0.9)	(0.7)		(0.9)		
急速進行性糸球体腎炎	697	715	590	271	121	85	26	19	6	5	2,535	3		2,538	5.94	6.59
(%)	(1.0)	(0.9)	(0.7)	(0.7)	(0.5)	(0.7)	(0.4)	(0.5)	(0.3)	(0.6)	(0.8)	(1.0)		(0.8)		
妊娠腎/妊娠中毒症	62	114	208	208	256	217	195	139	64	20	1,483			1,483	17.73	10.29
(%)	(0.1)	(0.1)	(0.3)	(0.5)	(1.2)	(1.7)	(2.8)	(3.5)	(3.0)	(2.5)	(0.5)			(0.5)		
その他の分類不能の腎炎	282	297	301	205	118	105	77	42	23	4	1,454			1,454	9.70	9.48
(%)	(0.4)	(0.4)	(0.4)	(0.5)	(0.5)	(0.8)	(1.1)	(1.1)	(1.1)	(0.5)	(0.5)			(0.5)		
多発性嚢胞腎	1,924	2,516	2,953	1,999	1,144	586	232	79	30	6	11,469	2		11,471	8.20	7.08
(%)	(2.7)	(3.1)	(3.7)	(4.9)	(5.2)	(4.7)	(3.4)	(2.0)	(1.4)	(0.8)	(3.6)	(0.7)		(3.6)		
腎硬化症	9,902	9,991	7,617	2,645	965	335	126	35	15	6	31,637	13		31,650	4.52	4.82
(%)	(14.0)	(12.5)	(9.6)	(6.5)	(4.4)	(2.7)	(1.8)	(0.9)	(0.7)	(0.8)	(9.9)	(4.4)		(9.9)		
悪性高血圧	604	681	716	351	219	94	40	11	5		2,721			2,721	6.83	6.53
(%)	(0.9)	(0.8)	(0.9)	(0.9)	(1.0)	(0.8)	(0.6)	(0.3)	(0.2)		(0.9)			(0.9)		
糖尿病性腎症	31,871	37,305	35,309	13,851	4,046	963	202	50	13	3	123,613	44		123,657	4.98	4.57
(%)	(45.2)	(46.5)	(44.6)	(34.2)	(18.3)	(7.7)	(2.9)	(1.3)	(0.6)	(0.4)	(38.8)	(14.9)		(38.8)		
SLE腎炎	418	475	510	304	239	129	77	45	21	2	2,220	2		2,222	8.92	8.41
(%)	(0.6)	(0.6)	(0.6)	(0.7)	(1.1)	(1.0)	(1.1)	(1.1)	(1.0)	(0.3)	(0.7)	(0.7)		(0.7)		
アミロイド腎	143	126	100	42	15	4	5	1	2		438			438	4.95	5.71
(%)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.0)	(0.1)	(0.0)	(0.1)		(0.1)			(0.1)		
痛風腎	153	201	264	178	128	78	34	12	5	2	1,055	1		1,056	9.49	7.81
(%)	(0.2)	(0.3)	(0.3)	(0.4)	(0.6)	(0.6)	(0.5)	(0.3)	(0.2)	(0.3)	(0.3)	(0.3)		(0.3)		
先天性代謝異常による腎不全	44	46	64	53	25	14	19	7	6	2	280	1		281	10.65	9.60
(%)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.1)	(0.3)	(0.2)	(0.3)	(0.3)	(0.1)	(0.3)		(0.1)		
腎・尿路結核	17	33	31	24	17	18	12	7	7	2	168			168	12.98	10.83
(%)	(0.0)	(0.0)	(0.0)	(0.1)	(0.1)	(0.1)	(0.2)	(0.2)	(0.3)	(0.3)	(0.1)			(0.1)		
腎・尿路結石	119	132	161	74	42	33	9	3	5		578			578	7.60	7.30
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.3)	(0.1)	(0.1)	(0.2)		(0.2)			(0.2)		
腎・尿路腫瘍	304	269	238	88	34	15	5	5	1		959			959	4.95	5.54
(%)	(0.4)	(0.3)	(0.3)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)	(0.0)		(0.3)			(0.3)		
閉塞性尿路障害	158	166	172	93	63	39	35	12	3	1	742	1		743	8.39	8.28
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.5)	(0.3)	(0.1)	(0.1)	(0.2)	(0.3)		(0.2)		
骨髄腫	173	86	48	11	2	1	1	1			323			323	2.60	3.87
(%)	(0.2)	(0.1)	(0.1)	(0.0)	(0.0)	(0.0)	(0.0)	(0.0)			(0.1)			(0.1)		
腎形成不全	91	115	121	98	81	71	43	25	13	6	664			664	11.97	9.93
(%)	(0.1)	(0.1)	(0.2)	(0.2)	(0.4)	(0.6)	(0.6)	(0.6)	(0.6)	(0.8)	(0.2)			(0.2)		
不明	8,289	8,106	7,395	3,666	1,965	910	449	257	148	63	31,248	164		31,412	6.43	6.97
(%)	(11.8)	(10.1)	(9.3)	(9.0)	(8.9)	(7.3)	(6.5)	(6.4)	(6.9)	(7.9)	(9.8)	(55.6)		(9.8)		
再導入	377	401	466	303	192	140	140	98	37	18	2,172	2		2,174	10.75	10.07
(%)	(0.5)	(0.5)	(0.6)	(0.7)	(0.9)	(1.1)	(2.0)	(2.5)	(1.7)	(2.3)	(0.7)	(0.7)		(0.7)		
その他	2,393	2,133	1,946	868	507	252	157	87	57	18	8,418	20		8,438	6.44	7.35
(%)	(3.4)	(2.7)	(2.5)	(2.1)	(2.3)	(2.0)	(2.3)	(2.2)	(2.6)	(2.3)	(2.6)	(6.8)		(2.6)		
合計	70,538	80,168	79,145	40,556	22,094	12,484	6,897	3,987	2,152	793	318,814	295		319,109	7.35	7.60
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)		
記載なし																
総計	70,538	80,168	79,145	40,556	22,094	12,484	6,897	3,987	2,152	793	318,814	295		319,109	7.35	7.60

数値下のかっこ内は列方向の合計に対する％です。

##A0154##

(C)Japanese Society for Dialysis Therapy

2016年調査