

表 220 2016年末患者，透析歴，原疾患別人数(男性)

原疾患	2年未満	2年～	5年～	10年～	15年～	20年～	25年～	30年～	35年～	40年～	合計	不明	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	7,722	9,976	12,011	8,745	6,686	4,644	2,662	1,603	903	333	55,285	24		55,309	10.91	9.35
(%)	(16.1)	(18.5)	(23.2)	(35.0)	(52.5)	(67.1)	(73.5)	(77.7)	(79.3)	(82.0)	(26.9)	(12.8)		(26.9)		
慢性腎盂腎炎	237	264	278	231	124	90	47	49	26	3	1,349	2		1,351	9.95	9.33
(%)	(0.5)	(0.5)	(0.5)	(0.9)	(1.0)	(1.3)	(1.3)	(2.4)	(2.3)	(0.7)	(0.7)	(1.1)		(0.7)		
急速進行性糸球体腎炎	371	381	293	130	56	39	10	12	4	4	1,300	1		1,301	5.75	6.63
(%)	(0.8)	(0.7)	(0.6)	(0.5)	(0.4)	(0.6)	(0.3)	(0.6)	(0.4)	(1.0)	(0.6)	(0.5)		(0.6)		
妊娠腎/妊娠中毒症																
(%)																
その他の分類不能の腎炎	177	173	165	126	70	69	50	18	17	4	869			869	9.88	9.71
(%)	(0.4)	(0.3)	(0.3)	(0.5)	(0.5)	(1.0)	(1.4)	(0.9)	(1.5)	(1.0)	(0.4)			(0.4)		
多発性嚢胞腎	1,086	1,400	1,666	1,068	562	305	125	33	22	3	6,270			6,270	7.97	7.00
(%)	(2.3)	(2.6)	(3.2)	(4.3)	(4.4)	(4.4)	(3.5)	(1.6)	(1.9)	(0.7)	(3.0)			(3.0)		
腎硬化症	6,760	6,708	4,987	1,695	613	205	74	24	9	2	21,077	8		21,085	4.40	4.71
(%)	(14.1)	(12.4)	(9.6)	(6.8)	(4.8)	(3.0)	(2.0)	(1.2)	(0.8)	(0.5)	(10.2)	(4.3)		(10.2)		
悪性高血圧	414	471	488	239	154	52	21	5	4		1,848			1,848	6.67	6.33
(%)	(0.9)	(0.9)	(0.9)	(1.0)	(1.2)	(0.8)	(0.6)	(0.2)	(0.4)		(0.9)			(0.9)		
糖尿病性腎症	23,147	26,840	24,843	9,332	2,550	567	99	23	7	1	87,409	32		87,441	4.81	4.41
(%)	(48.2)	(49.7)	(47.9)	(37.3)	(20.0)	(8.2)	(2.7)	(1.1)	(0.6)	(0.2)	(42.5)	(17.0)		(42.5)		
SLE腎炎	159	180	166	95	66	28	13	4	2		713			713	7.06	6.95
(%)	(0.3)	(0.3)	(0.3)	(0.4)	(0.5)	(0.4)	(0.4)	(0.2)	(0.2)		(0.3)			(0.3)		
アミロイド腎	58	58	46	6	7		3	1	2		181			181	4.88	6.43
(%)	(0.1)	(0.1)	(0.1)	(0.0)	(0.1)		(0.1)	(0.0)	(0.2)		(0.1)			(0.1)		
痛風腎	126	185	230	154	117	64	30	9	5	2	922	1		923	9.50	7.80
(%)	(0.3)	(0.3)	(0.4)	(0.6)	(0.9)	(0.9)	(0.8)	(0.4)	(0.4)	(0.5)	(0.4)	(0.5)		(0.4)		
先天性代謝異常による腎不全	26	27	38	32	17	11	11	3	4	1	170			170	10.84	9.47
(%)	(0.1)	(0.0)	(0.1)	(0.1)	(0.1)	(0.2)	(0.3)	(0.1)	(0.4)	(0.2)	(0.1)			(0.1)		
腎・尿路結核	11	21	13	13	8	7	8	2	1	1	85			85	11.33	10.16
(%)	(0.0)	(0.0)	(0.0)	(0.1)	(0.1)	(0.1)	(0.2)	(0.1)	(0.1)	(0.2)	(0.0)			(0.0)		
腎・尿路結石	80	77	112	52	31	20	4	2	3		381			381	7.62	7.08
(%)	(0.2)	(0.1)	(0.2)	(0.2)	(0.2)	(0.3)	(0.1)	(0.1)	(0.3)		(0.2)			(0.2)		
腎・尿路腫瘍	248	215	180	57	23	11	3	4	1		742			742	4.67	5.41
(%)	(0.5)	(0.4)	(0.3)	(0.2)	(0.2)	(0.2)	(0.1)	(0.2)	(0.1)		(0.4)			(0.4)		
閉塞性尿路障害	111	119	117	52	36	22	22	9	1		489	1		490	7.81	7.97
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.3)	(0.3)	(0.6)	(0.4)	(0.1)		(0.2)	(0.5)		(0.2)		
骨髄腫	99	54	30	7	1		1				192			192	2.60	3.55
(%)	(0.2)	(0.1)	(0.1)	(0.0)	(0.0)		(0.0)				(0.1)			(0.1)		
腎形成不全	58	71	79	62	54	40	24	16	9	1	414			414	11.63	9.67
(%)	(0.1)	(0.1)	(0.2)	(0.2)	(0.4)	(0.6)	(0.7)	(0.8)	(0.8)	(0.2)	(0.2)			(0.2)		
不明	5,336	5,171	4,612	2,185	1,131	505	245	127	61	35	19,408	102		19,510	6.08	6.62
(%)	(11.1)	(9.6)	(8.9)	(8.7)	(8.9)	(7.3)	(6.8)	(6.2)	(5.4)	(8.6)	(9.4)	(54.3)		(9.5)		
再導入	236	258	304	198	132	95	97	70	29	7	1,426	2		1,428	11.00	10.11
(%)	(0.5)	(0.5)	(0.6)	(0.8)	(1.0)	(1.4)	(2.7)	(3.4)	(2.5)	(1.7)	(0.7)	(1.1)		(0.7)		
その他	1,529	1,379	1,172	522	299	146	72	48	29	9	5,205	15		5,220	6.06	7.00
(%)	(3.2)	(2.6)	(2.3)	(2.1)	(2.3)	(2.1)	(2.0)	(2.3)	(2.5)	(2.2)	(2.5)	(8.0)		(2.5)		
合計	47,991	54,028	51,830	25,001	12,737	6,920	3,621	2,062	1,139	406	205,735	188		205,923	6.84	7.19
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)		
記載なし																
総計	47,991	54,028	51,830	25,001	12,737	6,920	3,621	2,062	1,139	406	205,735	188		205,923	6.84	7.19

数値下のかっこ内は列方向の合計に対する%です。

##A0155##

(C)Japanese Society for Dialysis Therapy

2016年調査