

表 224 2016年末患者, 治療方法 (新分類), 性別

治療方法 (新分類)	男性	女性	合計	記載なし	総計
施設血液透析	150,885	82,818	233,703	1	233,704
(%)	(73.3)	(73.2)	(73.2)	(100.0)	(73.2)
Off-line HDF	2,817	1,820	4,637		4,637
(%)	(1.4)	(1.6)	(1.5)		(1.5)
On-line HDF	38,550	20,566	59,116		59,116
(%)	(18.7)	(18.2)	(18.5)		(18.5)
Push/Pull HDF	43	23	66		66
(%)	(0.0)	(0.0)	(0.0)		(0.0)
ハイオフィльтраーション	138	114	252		252
(%)	(0.1)	(0.1)	(0.1)		(0.1)
IHDF	6,743	3,985	10,728		10,728
(%)	(3.3)	(3.5)	(3.4)		(3.4)
血液濾過	13	6	19		19
(%)	(0.0)	(0.0)	(0.0)		(0.0)
血液吸着透析 (リクセル等使用)	662	693	1,355		1,355
(%)	(0.3)	(0.6)	(0.4)		(0.4)
在宅血液透析	470	147	617		617
(%)	(0.2)	(0.1)	(0.2)		(0.2)
腹膜透析 (CAPD)	3,110	1,863	4,973		4,973
(%)	(1.5)	(1.6)	(1.6)		(1.6)
腹膜透析 (APD)	1,765	872	2,637		2,637
(%)	(0.9)	(0.8)	(0.8)		(0.8)
腹膜透析 (CCPD)	727	278	1,005		1,005
(%)	(0.4)	(0.2)	(0.3)		(0.3)
合計	205,923	113,185	319,108	1	319,109
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
記載なし					
総計	205,923	113,185	319,108	1	319,109

数値下のかっこ内は列方向の合計に対する%です。

##A0159##

(C)Japanese Society for Dialysis Therapy

2016年調査