

表 244 2016年死亡患者(透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1年～	2年～	5年～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	193	155	259	417	1,152	1,532	1,022	702	483	820	6,735	48		6,783
(%)	(13.6)	(13.9)	(15.6)	(15.7)	(16.2)	(18.6)	(24.8)	(37.9)	(53.6)	(66.2)	(22.2)	(15.3)		(22.1)
慢性腎盂腎炎	11	7	10	19	48	43	26	21	11	25	221	2		223
(%)	(0.8)	(0.6)	(0.6)	(0.7)	(0.7)	(0.5)	(0.6)	(1.1)	(1.2)	(2.0)	(0.7)	(0.6)		(0.7)
急速進行性糸球体腎炎	47	38	32	33	75	60	24	6	4	2	321	4		325
(%)	(3.3)	(3.4)	(1.9)	(1.2)	(1.1)	(0.7)	(0.6)	(0.3)	(0.4)	(0.2)	(1.1)	(1.3)		(1.1)
妊娠腎/妊娠中毒症					2	5	7	9	11	27	61	1		62
(%)					(0.0)	(0.1)	(0.2)	(0.5)	(1.2)	(2.2)	(0.2)	(0.3)		(0.2)
その他の分類不能の腎炎	7	7	8	7	23	20	12	6	8	8	106			106
(%)	(0.5)	(0.6)	(0.5)	(0.3)	(0.3)	(0.2)	(0.3)	(0.3)	(0.9)	(0.6)	(0.3)			(0.3)
多発性嚢胞腎	13	10	12	35	117	154	119	90	45	45	640	5		645
(%)	(0.9)	(0.9)	(0.7)	(1.3)	(1.6)	(1.9)	(2.9)	(4.9)	(5.0)	(3.6)	(2.1)	(1.6)		(2.1)
腎硬化症	189	140	225	440	1,102	925	355	106	36	26	3,544	17		3,561
(%)	(13.3)	(12.6)	(13.6)	(16.5)	(15.5)	(11.2)	(8.6)	(5.7)	(4.0)	(2.1)	(11.7)	(5.4)		(11.6)
悪性高血圧	9	2	4	19	32	48	23	11	9	8	165	2		167
(%)	(0.6)	(0.2)	(0.2)	(0.7)	(0.4)	(0.6)	(0.6)	(0.6)	(1.0)	(0.6)	(0.5)	(0.6)		(0.5)
糖尿病性腎症	356	341	622	1,045	3,205	4,052	1,909	598	143	62	12,333	108		12,441
(%)	(25.0)	(30.7)	(37.5)	(39.3)	(45.0)	(49.2)	(46.3)	(32.3)	(15.9)	(5.0)	(40.7)	(34.5)		(40.6)
SLE腎炎	12	14	11	18	44	43	24	13	5	16	200			200
(%)	(0.8)	(1.3)	(0.7)	(0.7)	(0.6)	(0.5)	(0.6)	(0.7)	(0.6)	(1.3)	(0.7)			(0.7)
アミロイド腎	7	17	11	6	21	19	3	2			86			86
(%)	(0.5)	(1.5)	(0.7)	(0.2)	(0.3)	(0.2)	(0.1)	(0.1)			(0.3)			(0.3)
痛風腎	2	2	3		7	10	16	9	9	9	67			67
(%)	(0.1)	(0.2)	(0.2)		(0.1)	(0.1)	(0.4)	(0.5)	(1.0)	(0.7)	(0.2)			(0.2)
先天性代謝異常による腎不全	1	1		1	3	3	1	3	1	1	15			15
(%)	(0.1)	(0.1)		(0.0)	(0.0)	(0.0)	(0.0)	(0.2)	(0.1)	(0.1)	(0.0)			(0.0)
腎・尿路結核					4	2	6	4	3	4	23			23
(%)					(0.1)	(0.0)	(0.1)	(0.2)	(0.3)	(0.3)	(0.1)			(0.1)
腎・尿路結石	3	2	4	6	10	15	8	3	1	5	57			57
(%)	(0.2)	(0.2)	(0.2)	(0.2)	(0.1)	(0.2)	(0.2)	(0.2)	(0.1)	(0.4)	(0.2)			(0.2)
腎・尿路腫瘍	11	7	23	26	30	19	11	4	1	2	134	2		136
(%)	(0.8)	(0.6)	(1.4)	(1.0)	(0.4)	(0.2)	(0.3)	(0.2)	(0.1)	(0.2)	(0.4)	(0.6)		(0.4)
閉塞性尿路障害	11	5	4	6	17	21	6	5	1	2	78			78
(%)	(0.8)	(0.4)	(0.2)	(0.2)	(0.2)	(0.3)	(0.1)	(0.3)	(0.1)	(0.2)	(0.3)			(0.3)
骨髄腫	19	22	21	21	26	7					116			116
(%)	(1.3)	(2.0)	(1.3)	(0.8)	(0.4)	(0.1)					(0.4)			(0.4)
腎形成不全	1	1		2	6	2	2	2			16			16
(%)	(0.1)	(0.1)		(0.1)	(0.1)	(0.0)	(0.0)	(0.1)			(0.1)			(0.1)
不明	346	236	289	413	970	987	437	211	112	144	4,145	108		4,253
(%)	(24.3)	(21.2)	(17.4)	(15.5)	(13.6)	(12.0)	(10.6)	(11.4)	(12.4)	(11.6)	(13.7)	(34.5)		(13.9)
再導入	13	9	6	12	15	32	11	8	3	11	120	2		122
(%)	(0.9)	(0.8)	(0.4)	(0.5)	(0.2)	(0.4)	(0.3)	(0.4)	(0.3)	(0.9)	(0.4)	(0.6)		(0.4)
その他	173	96	116	136	215	236	98	37	15	22	1,144	14		1,158
(%)	(12.1)	(8.6)	(7.0)	(5.1)	(3.0)	(2.9)	(2.4)	(2.0)	(1.7)	(1.8)	(3.8)	(4.5)		(3.8)
合計	1,424	1,112	1,660	2,662	7,124	8,235	4,120	1,850	901	1,239	30,327	313		30,640
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし														
総計	1,424	1,112	1,660	2,662	7,124	8,235	4,120	1,850	901	1,239	30,327	313		30,640

数値下のかっこ内は列方向の合計に対する%です。

##A0179##

(C)Japanese Society for Dialysis Therapy

2016年調査