

表 245 2016年死亡患者(男性 年齢、原疾患)

原疾患	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差						
慢性系球体腎炎 (%)				1 (25.0)	3 (50.0)	1 (11.1)	4 (18.2)	19 (22.9)	43 (18.6)	71 (19.2)	145 (22.1)	261 (20.7)	590 (21.2)	567 (19.8)	753 (20.0)	886 (21.6)	631 (22.0)	227 (22.5)	43 (23.9)	4,245 (21.0)	4,245 (21.0)	4,245 (21.0)	4,245 (21.0)	75.92 (21.0)	10.35 (21.0)							
慢性腎孟腎炎 (%)								1 (0.4)	8 (2.2)	7 (1.1)	11 (0.9)	13 (0.5)	21 (0.7)	16 (0.4)	21 (0.5)	13 (0.5)	7 (0.7)	13 (0.5)	9 (0.5)	7 (0.7)	118 (0.6)	118 (0.6)	118 (0.6)	118 (0.6)	73.49 (0.6)	11.08 (0.6)						
急速進行性系球体腎炎 (%)										3 (0.5)	4 (0.3)	20 (0.7)	26 (0.9)	44 (1.2)	40 (1.0)	24 (0.8)	24 (0.9)	9 (0.9)	9 (0.9)	7 (0.8)	7 (0.8)	170 (0.8)	170 (0.8)	170 (0.8)	170 (0.8)	77.82 (0.8)	7.73 (0.8)					
妊娠腎/妊娠中毒症 (%)																																
その他の分類不能の腎炎 (%)			1 (100.0)							1 (0.4)	1 (0.3)	1 (0.2)	14 (0.5)	11 (0.4)	16 (0.3)	13 (0.3)	10 (0.3)	1 (0.1)	3 (0.1)	72 (1.7)	72 (1.7)	72 (1.7)	72 (1.7)	75.82 (0.4)	11.62 (0.4)							
多発性囊胞腎 (%)										3 (3.6)	7 (3.0)	11 (3.0)	29 (2.4)	77 (2.8)	68 (2.4)	62 (1.6)	56 (1.4)	45 (1.6)	11 (1.1)	2 (1.1)	2 (1.1)	2 (1.1)	2 (1.1)	2 (1.1)	2 (1.1)	387 (1.9)	10.42 (1.9)					
腎硬化症 (%)										2 (33.3)	1 (11.1)	4 (18.2)	6 (7.2)	12 (5.2)	16 (4.3)	24 (3.7)	43 (3.4)	140 (5.0)	192 (6.7)	388 (10.3)	622 (15.2)	580 (20.2)	273 (27.1)	47 (27.1)	2,350 (26.1)	2,350 (26.1)	2,350 (26.1)	2,350 (26.1)	80.94 (11.6)	9.00 (11.6)		
悪性高血圧 (%)										1 (4.5)	1 (0.4)	1 (0.8)	3 (0.5)	5 (0.4)	7 (0.3)	16 (0.6)	19 (0.5)	20 (0.5)	17 (0.7)	1 (0.6)	1 (0.6)	1 (0.6)	1 (0.6)	1 (0.6)	1 (0.6)	113 (0.6)	113 (0.6)	113 (0.6)	113 (0.6)	78.16 (0.6)	11.22 (0.6)	
糖尿病性腎症 (%)	1 (100.0)							1 (25.0)	2 (22.2)	8 (36.4)	39 (47.0)	121 (52.4)	186 (50.8)	701 (56.2)	1,440 (55.5)	1,393 (51.7)	1,746 (48.7)	1,617 (46.5)	895 (39.5)	216 (31.2)	30 (21.4)	30 (16.7)	30 (16.7)	30 (16.7)	30 (16.7)	30 (16.7)	8,768 (14.3)	8,768 (14.3)	8,768 (14.3)	8,768 (14.3)	9.88 (14.3)	
SLE腎炎 (%)										1 (1.2)	1 (0.4)	1 (0.3)	3 (0.5)	5 (0.4)	12 (0.4)	14 (0.5)	11 (0.3)	16 (0.4)	6 (0.2)	1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	71 (0.4)	71 (0.4)	71 (0.4)	71 (0.4)	73.10 (0.4)	10.21 (0.4)	
アミロイド腎 (%)										3 (3.6)	1 (0.4)	1 (0.2)	5 (0.3)	6 (0.2)	10 (0.2)	10 (0.3)	8 (0.2)	2 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	41 (0.2)	41 (0.2)	41 (0.2)	41 (0.2)	71.73 (0.2)	12.27 (0.2)			
痛風腎 (%)											1 (0.3)	1 (0.1)	1 (0.1)	7 (0.4)	12 (0.4)	14 (0.4)	12 (0.4)	10 (0.4)	2 (0.2)	59 (0.3)	59 (0.3)	59 (0.3)	59 (0.3)	59 (0.3)	59 (0.3)	59 (0.3)	59 (0.3)	59 (0.3)	59 (0.3)	77.46 (0.3)	7.95 (0.3)	
先天性代謝異常による腎不全 (%)																					2 (0.3)	2 (0.3)	2 (0.3)	2 (0.3)	2 (0.3)	2 (0.3)	2 (0.3)	2 (0.3)	2 (0.3)	2 (0.3)	5 (0.0)	14.54 (0.0)
腎・尿路結核 (%)																																
腎・尿路結石 (%)																																
腎・尿路腫瘍 (%)																																
閉塞性尿路障害 (%)																																
骨髄腫 (%)																																
腎形成不全 (%)	1 (100.0)																															
不明 (%)																																
再導入 (%)																																
その他 (%)																																
合計 (%)	1 (100.0)	1 (100.0)	1 (100.0)	1 (100.0)	4 (100.0)	6 (100.0)	9 (100.0)	22 (100.0)	83 (100.0)	231 (100.0)	370 (100.0)	655 (100.0)	1,263 (100.0)	2,787 (100.0)	2,856 (100.0)	3,758 (100.0)	4,093 (100.0)	2,868 (100.0)	2,868 (100.0)	1,008 (100.0)	180 (100.0)	20,199 (100.0)	20,199 (100.0)	20,199 (100.0)	20,199 (100.0)	75.68 (10.0)	10.28 (10.0)					
記載なし																																
総計	1	1	1	1	1	4	6	9	22	83	231	370	655	1,263	2,787	2,856	3,758	4,093	2,868	1,008	180	20,199	20,199	20,199	20,199	75.68 (10.0)	10.28 (10.0)					

数値下のかっこ内は列方向の合計に対する%です。

##A0180##

(C)Japanese Society for Dialysis Therapy

2016年調査