

表 249 2016年死亡患者(男性 透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1年～	2年～	5年～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	117	94	178	285	773	957	640	435	279	457	4,215	30		4,245
(%)	(12.5)	(12.5)	(15.8)	(15.3)	(15.8)	(17.5)	(24.6)	(38.0)	(53.2)	(67.3)	(21.1)	(14.2)		(21.0)
慢性腎孟腎炎	8	5	6	11	25	22	11	9	5	14	116	2		118
(%)	(0.9)	(0.7)	(0.5)	(0.6)	(0.5)	(0.4)	(0.4)	(0.8)	(1.0)	(2.1)	(0.6)	(0.9)		(0.6)
急速進行性糸球体腎炎	21	23	23	13	40	29	14	2	2	1	168	2		170
(%)	(2.2)	(3.1)	(2.0)	(0.7)	(0.8)	(0.5)	(0.5)	(0.2)	(0.4)	(0.1)	(0.8)	(0.9)		(0.8)
妊娠腎/妊娠中毒症														
(%)														
その他の分類不能の腎炎	4	6	6	6	13	13	8	5	5	6	72			72
(%)	(0.4)	(0.8)	(0.5)	(0.3)	(0.3)	(0.2)	(0.3)	(0.4)	(1.0)	(0.9)	(0.4)			(0.4)
多発性囊胞腎	8	2	7	23	78	96	71	43	31	25	384	3		387
(%)	(0.9)	(0.3)	(0.6)	(1.2)	(1.6)	(1.8)	(2.7)	(3.8)	(5.9)	(3.7)	(1.9)	(1.4)		(1.9)
腎硬化症	135	98	160	289	743	590	218	72	19	19	2,343	7		2,350
(%)	(14.4)	(13.0)	(14.2)	(15.5)	(15.2)	(10.8)	(8.4)	(6.3)	(3.6)	(2.8)	(11.7)	(3.3)		(11.6)
悪性高血圧	6	2	4	15	20	31	13	9	7	4	111	2		113
(%)	(0.6)	(0.3)	(0.4)	(0.8)	(0.4)	(0.6)	(0.5)	(0.8)	(1.3)	(0.6)	(0.6)	(0.9)		(0.6)
糖尿病性腎症	256	249	455	781	2,315	2,871	1,260	377	87	41	8,692	76		8,768
(%)	(27.4)	(33.1)	(40.4)	(41.9)	(47.4)	(52.4)	(48.5)	(32.9)	(16.6)	(6.0)	(43.5)	(36.0)		(43.4)
SLE腎炎	5	7	4	9	13	16	10	3		4	71			71
(%)	(0.5)	(0.9)	(0.4)	(0.5)	(0.3)	(0.3)	(0.4)	(0.3)		(0.6)	(0.4)			(0.4)
アミロイド腎	4	8	8	3	9	6	2	1			41			41
(%)	(0.4)	(1.1)	(0.7)	(0.2)	(0.2)	(0.1)	(0.1)	(0.1)			(0.2)			(0.2)
痛風腎	2	2	2		7	9	14	8	7	8	59			59
(%)	(0.2)	(0.3)	(0.2)		(0.1)	(0.2)	(0.5)	(0.7)	(1.3)	(1.2)	(0.3)			(0.3)
先天性代謝異常による腎不全	1			1	2			1			5			5
(%)	(0.1)			(0.1)	(0.0)			(0.1)			(0.0)			(0.0)
腎・尿路結核					3		4	3	3	2	15			15
(%)					(0.1)		(0.2)	(0.3)	(0.6)	(0.3)	(0.1)			(0.1)
腎・尿路結石	2	1	1	4	5	8	5	1		3	30			30
(%)	(0.2)	(0.1)	(0.1)	(0.2)	(0.1)	(0.1)	(0.2)	(0.1)		(0.4)	(0.2)			(0.1)
腎・尿路腫瘍	8	3	18	19	24	16	9	1	1	2	101	2		103
(%)	(0.9)	(0.4)	(1.6)	(1.0)	(0.5)	(0.3)	(0.3)	(0.1)	(0.2)	(0.3)	(0.5)	(0.9)		(0.5)
閉塞性尿路障害	6	4	2	3	12	12	4	4		1	48			48
(%)	(0.6)	(0.5)	(0.2)	(0.2)	(0.2)	(0.2)	(0.2)	(0.3)		(0.1)	(0.2)			(0.2)
骨髄腫	9	8	16	11	17	2					63			63
(%)	(1.0)	(1.1)	(1.4)	(0.6)	(0.3)	(0.0)					(0.3)			(0.3)
腎形成不全	1	1		2	3	1	1	1			10			10
(%)	(0.1)	(0.1)		(0.1)	(0.1)	(0.0)	(0.0)	(0.1)			(0.1)			(0.0)
不明	222	167	165	284	633	636	247	138	63	72	2,627	74		2,701
(%)	(23.7)	(22.2)	(14.6)	(15.2)	(13.0)	(11.6)	(9.5)	(12.0)	(10.6)	(13.1)	(35.1)			(13.4)
再導入	9	8	4	11	13	21	9	5	1	9	90	1		91
(%)	(1.0)	(1.1)	(0.4)	(0.6)	(0.3)	(0.4)	(0.3)	(0.4)	(0.2)	(1.3)	(0.5)	(0.5)		(0.5)
その他	112	64	68	93	132	147	58	28	14	11	727	12		739
(%)	(12.0)	(8.5)	(6.0)	(5.0)	(2.7)	(2.7)	(2.2)	(2.4)	(2.7)	(1.6)	(3.6)	(5.7)		(3.7)
合計	936	752	1,127	1,863	4,880	5,483	2,598	1,146	524	679	19,988	211		20,199
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし														
総計	936	752	1,127	1,863	4,880	5,483	2,598	1,146	524	679	19,988	211		20,199

数値下のかっこ内は列方向の合計に対する%です。

##A0184##

(C)Japanese Society for Dialysis Therapy

2016年調査