

表 250 2016年死亡患者(女性 透析歴、原疾患)

原疾患	3ヶ月未満	3ヶ月～	6ヶ月～	1年～	2年～	5年～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
慢性糸球体腎炎	76	61	81	132	379	575	382	267	204	363	2,520	18		2,538
(%)	(15.6)	(16.9)	(15.2)	(16.5)	(16.9)	(20.9)	(25.1)	(37.9)	(54.1)	(64.8)	(24.4)	(17.6)		(24.3)
慢性腎盂腎炎	3	2	4	8	23	21	15	12	6	11	105			105
(%)	(0.6)	(0.6)	(0.8)	(1.0)	(1.0)	(0.8)	(1.0)	(1.7)	(1.6)	(2.0)	(1.0)			(1.0)
急速進行性糸球体腎炎	26	15	9	20	35	31	10	4	2	1	153	2		155
(%)	(5.3)	(4.2)	(1.7)	(2.5)	(1.6)	(1.1)	(0.7)	(0.6)	(0.5)	(0.2)	(1.5)	(2.0)		(1.5)
妊娠腎/妊娠中毒症					2	5	7	9	11	27	61	1		62
(%)					(0.1)	(0.2)	(0.5)	(1.3)	(2.9)	(4.8)	(0.6)	(1.0)		(0.6)
その他の分類不能の腎炎	3	1	2	1	10	7	4	1	3	2	34			34
(%)	(0.6)	(0.3)	(0.4)	(0.1)	(0.4)	(0.3)	(0.3)	(0.1)	(0.8)	(0.4)	(0.3)			(0.3)
多発性嚢胞腎	5	8	5	12	39	58	48	47	14	20	256	2		258
(%)	(1.0)	(2.2)	(0.9)	(1.5)	(1.7)	(2.1)	(3.2)	(6.7)	(3.7)	(3.6)	(2.5)	(2.0)		(2.5)
腎硬化症	54	42	65	151	359	335	137	34	17	7	1,201	10		1,211
(%)	(11.1)	(11.7)	(12.2)	(18.9)	(16.0)	(12.2)	(9.0)	(4.8)	(4.5)	(1.3)	(11.6)	(9.8)		(11.6)
悪性高血圧	3			4	12	17	10	2	2	4	54			54
(%)	(0.6)			(0.5)	(0.5)	(0.6)	(0.7)	(0.3)	(0.5)	(0.7)	(0.5)			(0.5)
糖尿病性腎症	100	92	167	264	890	1,181	649	221	56	21	3,641	32		3,673
(%)	(20.5)	(25.6)	(31.3)	(33.0)	(39.7)	(42.9)	(42.6)	(31.4)	(14.9)	(3.8)	(35.2)	(31.4)		(35.2)
SLE腎炎	7	7	7	9	31	27	14	10	5	12	129			129
(%)	(1.4)	(1.9)	(1.3)	(1.1)	(1.4)	(1.0)	(0.9)	(1.4)	(1.3)	(2.1)	(1.2)			(1.2)
アミロイド腎	3	9	3	3	12	13	1	1			45			45
(%)	(0.6)	(2.5)	(0.6)	(0.4)	(0.5)	(0.5)	(0.1)	(0.1)			(0.4)			(0.4)
痛風腎			1			1	2	1	2	1	8			8
(%)			(0.2)			(0.0)	(0.1)	(0.1)	(0.5)	(0.2)	(0.1)			(0.1)
先天性代謝異常による腎不全		1			1	3	1	2	1	1	10			10
(%)		(0.3)			(0.0)	(0.1)	(0.1)	(0.3)	(0.3)	(0.2)	(0.1)			(0.1)
腎・尿路結核					1	2	2	1		2	8			8
(%)					(0.0)	(0.1)	(0.1)	(0.1)		(0.4)	(0.1)			(0.1)
腎・尿路結石	1	1	3	2	5	7	3	2	1	2	27			27
(%)	(0.2)	(0.3)	(0.6)	(0.3)	(0.2)	(0.3)	(0.2)	(0.3)	(0.3)	(0.4)	(0.3)			(0.3)
腎・尿路腫瘍	3	4	5	7	6	3	2	3			33			33
(%)	(0.6)	(1.1)	(0.9)	(0.9)	(0.3)	(0.1)	(0.1)	(0.4)			(0.3)			(0.3)
閉塞性尿路障害	5	1	2	3	5	9	2	1	1	1	30			30
(%)	(1.0)	(0.3)	(0.4)	(0.4)	(0.2)	(0.3)	(0.1)	(0.1)	(0.3)	(0.2)	(0.3)			(0.3)
骨髄腫	10	14	5	10	9	5					53			53
(%)	(2.0)	(3.9)	(0.9)	(1.3)	(0.4)	(0.2)					(0.5)			(0.5)
腎形成不全					3	1	1	1			6			6
(%)					(0.1)	(0.0)	(0.1)	(0.1)			(0.1)			(0.1)
不明	124	69	124	129	337	351	190	73	49	72	1,518	34		1,552
(%)	(25.4)	(19.2)	(23.3)	(16.1)	(15.0)	(12.8)	(12.5)	(10.4)	(13.0)	(12.9)	(14.7)	(33.3)		(14.9)
再導入	4	1	2	1	2	11	2	3	2	2	30	1		31
(%)	(0.8)	(0.3)	(0.4)	(0.1)	(0.1)	(0.4)	(0.1)	(0.4)	(0.5)	(0.4)	(0.3)	(1.0)		(0.3)
その他	61	32	48	43	83	89	40	9	1	11	417	2		419
(%)	(12.5)	(8.9)	(9.0)	(5.4)	(3.7)	(3.2)	(2.6)	(1.3)	(0.3)	(2.0)	(4.0)	(2.0)		(4.0)
合計	488	360	533	799	2,244	2,752	1,522	704	377	560	10,339	102		10,441
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)		(100.0)
記載なし														
総計	488	360	533	799	2,244	2,752	1,522	704	377	560	10,339	102		10,441

数値下のかっこ内は列方向の合計に対する%です。

##A0185##

(C)Japanese Society for Dialysis Therapy

2016年調査