

表 301 2016年死亡患者死亡原因分類(原疾患との関係)

原疾患	心不全	脳血管障害	感染症	消化管出血	悪性腫瘍	悪液質 /尿毒症	心筋梗塞	カリウム 中毒/頓死	肝硬変症	自殺/拒否	腸閉塞	肺血栓 /肺塞栓	災害 ・事故死	その他	不明	合計	記載なし	総計
慢性糸球体腎炎	1,842	432	1,431	89	680	275	238	161	62	49	76	19	44	689	696	6,783		6,783
(%)	(27.2)	(6.4)	(21.1)	(1.3)	(10.0)	(4.1)	(3.5)	(2.4)	(0.9)	(0.7)	(1.1)	(0.3)	(0.6)	(10.2)	(10.3)	(100.0)		
慢性腎盂腎炎	61	9	51	1	36	4	6	8	4	2	1	1	2	16	21	223		223
(%)	(27.4)	(4.0)	(22.9)	(0.4)	(16.1)	(1.8)	(2.7)	(3.6)	(1.8)	(0.9)	(0.4)	(0.4)	(0.9)	(7.2)	(9.4)	(100.0)		
急速進行性糸球体腎炎	70	22	110	8	27	17	7	5	4	2	3	1		27	22	325		325
(%)	(21.5)	(6.8)	(33.8)	(2.5)	(8.3)	(5.2)	(2.2)	(1.5)	(1.2)	(0.6)	(0.9)	(0.3)		(8.3)	(6.8)	(100.0)		
妊娠腎/妊娠中毒症	19	4	9	2	6	1	2				1		1	6	11	62		62
(%)	(30.6)	(6.5)	(14.5)	(3.2)	(9.7)	(1.6)	(3.2)				(1.6)		(1.6)	(9.7)	(17.7)	(100.0)		
その他の分類不能の腎炎	28	4	25		9	4	2		2	2	4			13	13	106		106
(%)	(26.4)	(3.8)	(23.6)		(8.5)	(3.8)	(1.9)		(1.9)	(1.9)	(3.8)			(12.3)	(12.3)	(100.0)		
多発性嚢胞腎	113	60	173	7	66	33	25	9	8	6	7		4	72	62	645		645
(%)	(17.5)	(9.3)	(26.8)	(1.1)	(10.2)	(5.1)	(3.9)	(1.4)	(1.2)	(0.9)	(1.1)		(0.6)	(11.2)	(9.6)	(100.0)		
腎硬化症	917	227	858	53	323	168	134	82	20	26	39	1	15	346	350	3,559	2	3,561
(%)	(25.8)	(6.4)	(24.1)	(1.5)	(9.1)	(4.7)	(3.8)	(2.3)	(0.6)	(0.7)	(1.1)	(0.0)	(0.4)	(9.7)	(9.8)	(100.0)		
悪性高血圧	47	18	28	2	14	11	3	5	2	2	3			18	14	167		167
(%)	(28.1)	(10.8)	(16.8)	(1.2)	(8.4)	(6.6)	(1.8)	(3.0)	(1.2)	(1.2)	(1.8)			(10.8)	(8.4)	(100.0)		
糖尿病性腎症	3,297	895	2,631	133	1,056	492	610	366	82	74	119	18	75	1,160	1,433	12,441		12,441
(%)	(26.5)	(7.2)	(21.1)	(1.1)	(8.5)	(4.0)	(4.9)	(2.9)	(0.7)	(0.6)	(1.0)	(0.1)	(0.6)	(9.3)	(11.5)	(100.0)		
SLE腎炎	57	7	56	2	13	9	5	4			3			29	15	200		200
(%)	(28.5)	(3.5)	(28.0)	(1.0)	(6.5)	(4.5)	(2.5)	(2.0)			(1.5)			(14.5)	(7.5)	(100.0)		
アミロイド腎	25	3	17		15	5		2	1	2				10	6	86		86
(%)	(29.1)	(3.5)	(19.8)		(17.4)	(5.8)		(2.3)	(1.2)	(2.3)				(11.6)	(7.0)	(100.0)		
痛風腎	12	3	21	2	9	3	1							5	11	67		67
(%)	(17.9)	(4.5)	(31.3)	(3.0)	(13.4)	(4.5)	(1.5)							(7.5)	(16.4)	(100.0)		
先天性代謝異常による腎不全	4	2	2	1	2	1							1	2		15		15
(%)	(26.7)	(13.3)	(13.3)	(6.7)	(13.3)	(6.7)							(6.7)	(13.3)		(100.0)		
腎・尿路結核	6	1	7		3	1							1	2	2	23		23
(%)	(26.1)	(4.3)	(30.4)		(13.0)	(4.3)							(4.3)	(8.7)	(8.7)	(100.0)		
腎・尿路結石	14	2	7	2	11	2	1	2	3		1			7	5	57		57
(%)	(24.6)	(3.5)	(12.3)	(3.5)	(19.3)	(3.5)	(1.8)	(3.5)	(5.3)		(1.8)			(12.3)	(8.8)	(100.0)		
腎・尿路腫瘍	13	2	22	2	72	1	2	1				2		8	11	136		136
(%)	(9.6)	(1.5)	(16.2)	(1.5)	(52.9)	(0.7)	(1.5)	(0.7)				(1.5)		(5.9)	(8.1)	(100.0)		
閉塞性尿路障害	17	1	12	1	18	5	2	1		1	2		2	7	9	78		78
(%)	(21.8)	(1.3)	(15.4)	(1.3)	(23.1)	(6.4)	(2.6)	(1.3)		(1.3)	(2.6)		(2.6)	(9.0)	(11.5)	(100.0)		
骨髄腫	13	3	13	2	60	5		1		1		2		4	12	116		116
(%)	(11.2)	(2.6)	(11.2)	(1.7)	(51.7)	(4.3)		(0.9)		(0.9)		(1.7)		(3.4)	(10.3)	(100.0)		
腎形成不全	6	2	1				1				1			2	3	16		16
(%)	(37.5)	(12.5)	(6.3)				(6.3)				(6.3)			(12.5)	(18.8)	(100.0)		
不明	1,026	228	950	66	406	215	137	77	47	22	53	11	13	459	543	4,253		4,253
(%)	(24.1)	(5.4)	(22.3)	(1.6)	(9.5)	(5.1)	(3.2)	(1.8)	(1.1)	(0.5)	(1.2)	(0.3)	(0.3)	(10.8)	(12.8)	(100.0)		
再導入	39	4	41	1	7	2	6				1			7	14	122		122
(%)	(32.0)	(3.3)	(33.6)	(0.8)	(5.7)	(1.6)	(4.9)				(0.8)			(5.7)	(11.5)	(100.0)		
その他	261	50	260	21	134	25	28	18	15	10	16	5	4	189	122	1,158		1,158
(%)	(22.5)	(4.3)	(22.5)	(1.8)	(11.6)	(2.2)	(2.4)	(1.6)	(1.3)	(0.9)	(1.4)	(0.4)	(0.3)	(16.3)	(10.5)	(100.0)		
合計	7,887	1,979	6,725	395	2,967	1,279	1,210	742	250	199	330	60	162	3,078	3,375	30,638	2	30,640
(%)	(25.7)	(6.5)	(21.9)	(1.3)	(9.7)	(4.2)	(3.9)	(2.4)	(0.8)	(0.6)	(1.1)	(0.2)	(0.5)	(10.0)	(11.0)	(100.0)		
記載なし																		
(%)																		
総計	7,887	1,979	6,725	395	2,967	1,279	1,210	742	250	199	330	60	162	3,078	3,375	30,638	2	30,640
(%)	(25.7)	(6.5)	(21.9)	(1.3)	(9.7)	(4.2)	(3.9)	(2.4)	(0.8)	(0.6)	(1.1)	(0.2)	(0.5)	(10.0)	(11.0)	(100.0)		

数値下のかっこ内は行方向の合計に対する%です。

##A0236##

(C)Japanese Society for Dialysis Therapy

2016年調査