

表 309 2016年死亡患者死原因分類(年齢別、原疾患別)原疾患 = 惠性高血圧

死因	5歳未満	5歳~	10歳~	15歳~	20歳~	25歳~	30歳~	35歳~	40歳~	45歳~	50歳~	55歳~	60歳~	65歳~	70歳~	75歳~	80歳~	85歳~	90歳~	95歳~	合計	不明	記載なし	総計	平均	標準偏差		
心不全 (%)																					47			47	78.60	8.54		
脳血管障害 (%)																					(28.1)			(28.1)	68.39	16.28		
感染症 (%)																					(108)			(108)	28	75.75	12.30	
消化管出血 (%)																					(16.8)			(16.8)	2	81.50	12.02	
悪性腫瘍 (%)																					(1.2)			(1.2)	14	79.93	7.64	
悪液質/尿毒症 (%)																					(8.4)			(8.4)	11	83.82	7.97	
心筋梗塞 (%)																					(6.6)			(6.6)	3	70.67	19.60	
カリウム中毒/頻死 (%)																					(1.8)			(1.8)	5	71.00	16.84	
肝硬変症 (%)																					(3.0)			(3.0)	2	74.00	2.83	
自殺/拒否 (%)																					(1.2)			(1.2)	2	81.50	0.71	
腸閉塞 (%)																					(1.2)			(1.2)	3	79.33	4.04	
肺血栓/肺塞栓 (%)																					(1.8)			(1.8)				
災害・事故死 (%)																												
その他 (%)																												
不明 (%)																												
合計 (%)								1	3	5	6	8	13	21	29	31	26	21	3	167				167	77.29	11.62		
記載なし																												
総計																												

数値下のかっこ内は列方向の合計に対する%です。

##A024##

(C)Japanese Society for Dialysis Therapy

2016年調査