

表 320 2016年死亡患者死亡原因分類(年齢別、原疾患別)原疾患=腎形成不全

死因	5歳未満	5歳～	10歳～	15歳～	20歳～	25歳～	30歳～	35歳～	40歳～	45歳～	50歳～	55歳～	60歳～	65歳～	70歳～	75歳～	80歳～	85歳～	90歳～	95歳～	合計	不明	記載なし	総計	平均	標準偏差
心不全 (%)				1 (100.0)											1 (50.0)	1 (100.0)	1 (100.0)	1 (33.3)	1 (100.0)	1 (37.5)	6	6 (37.5)	6 (37.5)	73.83	28.93	
脳血管障害 (%)										1 (100.0)										2	2 (12.5)	2 (12.5)	61.00	22.63		
感染症 (%)																				1 (6.3)	1 (6.3)	1 (6.3)	91.00			
消化管出血 (%)																										
悪性腫瘍 (%)																										
悪液質/尿毒症 (%)																										
心筋梗塞 (%)															1 (50.0)					1 (6.3)	1 (6.3)	1 (6.3)	60.00			
カリウム中毒/鎮死 (%)																										
肝硬変症 (%)																										
自殺/拒否 (%)																										
腸閉塞 (%)																				1 (6.3)	1 (6.3)	1 (6.3)	57.00			
肺血栓/肺塞栓 (%)																										
災害・事故死 (%)																										
その他 (%)	1 (100.0)							1 (100.0)											2 (12.5)	2 (12.5)	2 (12.5)	19.00	25.46			
不明 (%)																			3 (18.8)	3 (18.8)	3 (18.8)	75.33	14.64			
合計 (%)	1 (100.0)		1 (100.0)					1 (100.0)		1 (100.0)		2 (100.0)		2 (100.0)		1 (100.0)		1 (100.0)		1 (100.0)		16 (100.0)	16 (100.0)	64.81	27.80	
記載なし																										
総計	1		1					1		1		2		2		1		1		3		1		16	64.81	27.80

数値下のかっこ内は列方向の合計に対する%です。

##A025##

(C)Japanese Society for Dialysis Therapy

2016年調査