

表 324 2016年死亡患者死亡原因分類(治療方法別)

死因	施設血液透析	血液透析濾過	血液濾過	血液吸着透析	在宅血液透析	腹膜透析	離脱	生体腎移植	献腎移植	献腎移植 (生体腎か 献腎か不明 の場合)	合計	記載なし	総計
心不全	4,116	518	4	7	2	94					4,741	3,146	7,887
(%)	(26.8)	(26.2)	(57.1)	(20.6)	(28.6)	(22.9)					(26.6)	(24.5)	(25.7)
脳血管障害	951	141	1			21					1,114	865	1,979
(%)	(6.2)	(7.1)	(14.3)			(5.1)					(6.3)	(6.7)	(6.5)
感染症	3,329	368	1	5	2	86					3,791	2,934	6,725
(%)	(21.7)	(18.6)	(14.3)	(14.7)	(28.6)	(21.0)					(21.3)	(22.9)	(21.9)
消化管出血	186	20		1	1	6					214	181	395
(%)	(1.2)	(1.0)		(2.9)	(14.3)	(1.5)					(1.2)	(1.4)	(1.3)
悪性腫瘍	1,434	149		4	1	34					1,622	1,345	2,967
(%)	(9.3)	(7.5)		(11.8)	(14.3)	(8.3)					(9.1)	(10.5)	(9.7)
悪液質/尿毒症	615	61		4		29					709	570	1,279
(%)	(4.0)	(3.1)		(11.8)		(7.1)					(4.0)	(4.4)	(4.2)
心筋梗塞	617	88		1	1	16					723	487	1,210
(%)	(4.0)	(4.4)		(2.9)	(14.3)	(3.9)					(4.1)	(3.8)	(3.9)
カリウム中毒/頓死	364	76				12					452	290	742
(%)	(2.4)	(3.8)				(2.9)					(2.5)	(2.3)	(2.4)
肝硬変症	125	16		2		5					148	102	250
(%)	(0.8)	(0.8)		(5.9)		(1.2)					(0.8)	(0.8)	(0.8)
自殺/拒否	90	15				2					107	92	199
(%)	(0.6)	(0.8)				(0.5)					(0.6)	(0.7)	(0.6)
腸閉塞	161	19				7					187	143	330
(%)	(1.0)	(1.0)				(1.7)					(1.1)	(1.1)	(1.1)
肺血栓/肺塞栓	30	7									37	23	60
(%)	(0.2)	(0.4)									(0.2)	(0.2)	(0.2)
災害・事故死	76	17		1							94	68	162
(%)	(0.5)	(0.9)		(2.9)							(0.5)	(0.5)	(0.5)
その他	1,561	186		3		45					1,795	1,283	3,078
(%)	(10.2)	(9.4)		(8.8)		(11.0)					(10.1)	(10.0)	(10.0)
不明	1,714	297	1	6		53					2,071	1,304	3,375
(%)	(11.2)	(15.0)	(14.3)	(17.6)		(12.9)					(11.6)	(10.2)	(11.0)
合計	15,369	1,978	7	34	7	410					17,805	12,833	30,638
(%)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)					(100.0)	(100.0)	(100.0)
記載なし						2					2		2
総計	15,369	1,978	7	34	7	412					17,807	12,833	30,640

数値下のカッコ内は列方向の合計に対する%です。

##A0259##

(C)Japanese Society for Dialysis Therapy

2016年調査