

表 327 2016年死亡患者死亡原因の詳細(透析期間との関係)

死因	3ヶ月未満	3ヶ月～	6ヶ月～	1年～	2年～	5年～	10年～	15年～	20年～	25年～	合計	不明	記載なし	総計
心不全	262	207	332	576	1,498	1,824	892	399	199	240	6,429	65		6,494
(%)	(4.1)	(3.2)	(5.2)	(9.0)	(23.3)	(28.4)	(13.9)	(6.2)	(3.1)	(3.7)	(100.0)			
肺水腫(溢水)	22	17	14	14	83	71	54	13	2	8	298	3		301
(%)	(7.4)	(5.7)	(4.7)	(4.7)	(27.9)	(23.8)	(18.1)	(4.4)	(0.7)	(2.7)	(100.0)			
急性心筋梗塞(発症30日以内死亡)	38	19	33	71	196	238	118	56	25	26	820	8		828
(%)	(4.6)	(2.3)	(4.0)	(8.7)	(23.9)	(29.0)	(14.4)	(6.8)	(3.0)	(3.2)	(100.0)			
虚血性心疾患(急性心筋梗塞以外)	19	10	8	27	89	116	66	22	8	14	379	3		382
(%)	(5.0)	(2.6)	(2.1)	(7.1)	(23.5)	(30.6)	(17.4)	(5.8)	(2.1)	(3.7)	(100.0)			
不整脈、伝導障害	17	16	19	33	109	110	65	22	16	17	424	7		431
(%)	(4.0)	(3.8)	(4.5)	(7.8)	(25.7)	(25.9)	(15.3)	(5.2)	(3.8)	(4.0)	(100.0)			
心内膜炎および弁膜症	2	5	7	8	39	49	32	16	8	18	184	2		186
(%)	(1.1)	(2.7)	(3.8)	(4.3)	(21.2)	(26.6)	(17.4)	(8.7)	(4.3)	(9.8)	(100.0)			
その他の心疾患	26	18	28	44	105	125	59	30	16	22	473	2		475
(%)	(5.5)	(3.8)	(5.9)	(9.3)	(22.2)	(26.4)	(12.5)	(6.3)	(3.4)	(4.7)	(100.0)			
くも膜下出血	6	7	11	19	32	34	20	9	7	8	153	3		156
(%)	(3.9)	(4.6)	(7.2)	(12.4)	(20.9)	(22.2)	(13.1)	(5.9)	(4.6)	(5.2)	(100.0)			
脳内出血	38	26	45	60	229	273	119	60	39	27	916	8		924
(%)	(4.1)	(2.8)	(4.9)	(6.6)	(25.0)	(29.8)	(13.0)	(6.6)	(4.3)	(2.9)	(100.0)			
脳梗塞	31	22	41	56	160	201	99	36	24	21	691	11		702
(%)	(4.5)	(3.2)	(5.9)	(8.1)	(23.2)	(29.1)	(14.3)	(5.2)	(3.5)	(3.0)	(100.0)			
その他の脳血管疾患	6	7	8	18	51	60	24	14	3	5	196	1		197
(%)	(3.1)	(3.6)	(4.1)	(9.2)	(26.0)	(30.6)	(12.2)	(7.1)	(1.5)	(2.6)	(100.0)			
敗血症	176	92	119	183	448	659	357	178	93	143	2,448	28		2,476
(%)	(7.2)	(3.8)	(4.9)	(7.5)	(18.3)	(26.9)	(14.6)	(7.3)	(3.8)	(5.8)	(100.0)			
中枢神経系感染症	2		2	1	2	5	6		3		23			23
(%)	(8.7)		(8.7)	(4.3)	(8.7)	(21.7)	(26.1)		(13.0)		(100.0)			
肺炎	186	145	203	308	892	911	498	212	86	120	3,561	34		3,595
(%)	(5.2)	(4.1)	(5.7)	(8.6)	(25.0)	(25.6)	(14.0)	(6.0)	(2.4)	(3.4)	(100.0)			
インフルエンザ			1	2	6	15	1	2	1	3	31			31
(%)			(3.2)	(6.5)	(19.4)	(48.4)	(3.2)	(6.5)	(3.2)	(9.7)	(100.0)			
尿路感染症	2	1	3		5	8	2			3	24			24
(%)	(8.3)	(4.2)	(12.5)		(20.8)	(33.3)	(8.3)			(12.5)	(100.0)			
消化管・胆道系感染症・腹膜炎	10	7	11	17	56	50	45	13	7	11	227	1		228
(%)	(4.4)	(3.1)	(4.8)	(7.5)	(24.7)	(22.0)	(19.8)	(5.7)	(3.1)	(4.8)	(100.0)			
劇症(急性)ウイルス肝炎	1	1	1	1	4						8			8
(%)	(12.5)	(12.5)	(12.5)	(12.5)	(50.0)						(100.0)			
結核	5	8	2	5	8	7	2	3	1	2	43	1		44
(%)	(11.6)	(18.6)	(4.7)	(11.6)	(18.6)	(16.3)	(4.7)	(7.0)	(2.3)	(4.7)	(100.0)			
ヒト免疫不全ウイルス(HIV)感染症					1						1			1
(%)					(100.0)						(100.0)			
その他の感染症	17	12	14	24	61	74	42	23	7	18	292	3		295
(%)	(5.8)	(4.1)	(4.8)	(8.2)	(20.9)	(25.3)	(14.4)	(7.9)	(2.4)	(6.2)	(100.0)			
中枢神経系の悪性新生物	1		2	4	10	6	1	2	3	2	31			31
(%)	(3.2)		(6.5)	(12.9)	(32.3)	(19.4)	(3.2)	(6.5)	(9.7)	(6.5)	(100.0)			
呼吸器系の悪性新生物	11	18	29	70	156	141	49	32	8	16	530	5		535
(%)	(2.1)	(3.4)	(5.5)	(13.2)	(29.4)	(26.6)	(9.2)	(6.0)	(1.5)	(3.0)	(100.0)			
肝癌	17	21	30	54	115	112	38	16	5	15	423	5		428
(%)	(4.0)	(5.0)	(7.1)	(12.8)	(27.2)	(26.5)	(9.0)	(3.8)	(1.2)	(3.5)	(100.0)			
肝臓以外の消化器系の悪性新生物	36	35	61	92	195	216	75	38	17	27	792	2		794
(%)	(4.5)	(4.4)	(7.7)	(11.6)	(24.6)	(27.3)	(9.5)	(4.8)	(2.1)	(3.4)	(100.0)			
乳房の悪性新生物	1	1	5	5	12	6	9	4	3	5	51			51
(%)	(2.0)	(2.0)	(9.8)	(9.8)	(23.5)	(11.8)	(17.6)	(7.8)	(5.9)	(9.8)	(100.0)			
性器の悪性新生物	4	5	6	8	14	18	11	6	1	4	77			77
(%)	(5.2)	(6.5)	(7.8)	(10.4)	(18.2)	(23.4)	(14.3)	(7.8)	(1.3)	(5.2)	(100.0)			
腎の悪性新生物	9	15	20	31	37	33	20	16	20	35	236	1		237
(%)	(3.8)	(6.4)	(8.5)	(13.1)	(15.7)	(14.0)	(8.5)	(6.8)	(8.5)	(14.8)	(100.0)			
内分泌腺の悪性新生物	4	1	7	6	21	17	3	2	2	2	65			65
(%)	(6.2)	(1.5)	(10.8)	(9.2)	(32.3)	(26.2)	(4.6)	(3.1)	(3.1)	(3.1)	(100.0)			
造血・リンパ組織の悪性新生物	27	38	36	45	50	38	15	10	4	9	272	2		274
(%)	(9.9)	(14.0)	(13.2)	(16.5)	(18.4)	(14.0)	(5.5)	(3.7)	(1.5)	(3.3)	(100.0)			
その他の悪性新生物	32	22	40	64	104	115	42	19	15	18	471	4		475
(%)	(6.8)	(4.7)	(8.5)	(13.6)	(22.1)	(24.4)	(8.9)	(4.0)	(3.2)	(3.8)	(100.0)			
ウイルス性肝硬変	6	7	7	6	21	11	10	8	2	9	87	1		88
(%)	(6.9)	(8.0)	(8.0)	(6.9)	(24.1)	(12.6)	(11.5)	(9.2)	(2.3)	(10.3)	(100.0)			
ウイルス性以外の肝硬変	20	12	11	24	25	28	24	7	6	3	160	2		162
(%)	(12.5)	(7.5)	(6.9)	(15.0)	(15.6)	(17.5)	(15.0)	(4.4)	(3.8)	(1.9)	(100.0)			
腸の血行障害	7	9	8	8	44	41	38	16	6	14	191	1		192
(%)	(3.7)	(4.7)	(4.2)	(4.2)	(23.0)	(21.5)	(19.9)	(8.4)	(3.1)	(7.3)	(100.0)			
イレウス	11	10	2	12	31	23	16	11	4	12	132	2		134
(%)	(8.3)	(7.6)	(1.5)	(9.1)	(23.5)	(17.4)	(12.1)	(8.3)	(3.0)	(9.1)	(100.0)			
消化管出血	31	20	16	28	87	108	35	34	11	22	392	3		395
(%)	(7.9)	(5.1)	(4.1)	(7.1)	(22.2)	(27.6)	(8.9)	(8.7)	(2.8)	(5.6)	(100.0)			
被覆性腹膜硬化症					1	1			2		4			4
(%)					(25.0)	(25.0)			(50.0)		(100.0)			
その他の消化器疾患	13	7	19	23	51	41	33	13	4	19	223	2		225
(%)	(5.8)	(3.1)	(8.5)	(10.3)	(22.9)	(18.4)	(14.8)	(5.8)	(1.8)	(8.5)	(100.0)			
肺梗塞、肺塞栓症	3	3	4	5	12	20	9	2		2	60			60
(%)	(5.0)	(5.0)	(6.7)	(8.3)	(20.0)	(33.3)	(15.0)	(3.3)		(3.3)	(100.0)			
悪液質	11	25	33	51	165	199	112	32	13	37	678	6		684
(%)	(1.6)	(3.7)	(4.9)	(7.5)	(24.3)	(29.4)	(16.5)	(4.7)	(1.9)	(5.5)	(100.0)			
尿毒症	37	16	29	32	89	102	38	20	7	13	383	5		388
(%)	(9.7)	(4.2)	(7.6)	(8.4)	(23.2)	(26.6)	(9.9)	(5.2)	(1.8)	(3.4)	(100.0)			
認知症		2	2	3	22	24	3	1	1		58			58
(%)		(3.4)	(3.4)	(5.2)	(37.9)	(41.4)	(5.2)	(1.7)	(1.7)		(100.0)			
その他の悪液質/尿毒症	6	6	12	15	31	39	26	7	4	2	148	1		149
(%)	(4.1)	(4.1)	(8.1)	(10.1)	(20.9)	(26.4)	(17.6)	(4.7)	(2.7)	(1.4)	(100.0)			
高カリウム血症	3	3	3	5	26	20	11	7	1	2	81	2		83
(%)	(3.7)	(3.7)	(3.7)	(6.2)	(32.1)	(24.7)	(13.6)	(8.6)	(1.2)	(2.5)	(100.0)			
原因不明の突然死	10	18	35	54	168	197	99	43	12	19	655	4		659
(%)	(1.5)	(2.7)	(5.3)	(8.2)	(25.6)	(30.1)	(15.1)	(6.6)	(1.8)	(2.9)	(100.0)			
自殺	3		7	7	21	14	13	2	4	3	74			74
(%)	(4.1)		(9.5)	(9.5)	(28.4)	(18.9)	(17.6)	(2.7)	(5.4)	(4.1)	(100.0)			
治療拒否(透析拒否)	7	8	8	15	26	37	10	7	4	2	124	1		125
(%)	(5.6)	(6.5)	(6.5)	(12.1)	(21.0)	(29.8)	(8.1)	(5.6)	(3.2)	(1.6)	(100.0)			
災害・事故死	1	2	7	15	37	53	24	11	7	5	162			162
(%)	(0.6)	(1.2)	(4.3)	(9.3)	(22.8)	(32.7)	(14.8)	(6.8)	(4.3)	(3.1)	(100.0)			
その他	148	106	144	246	659	790	388	172	80	101	2,834	19		2,853
(%)	(5.2)	(3.7)	(5.1)	(8.7)	(23.3)	(27.9)	(13.7)	(6.1)	(2.8)	(3.6)	(100.0)			
不明	99	82	175	267	820	955	467	202	110	135	3,312	63		3,375
(%)	(3.0)	(2.5)	(5.3)	(8.1)	(24.8)	(28.8)	(14.1)	(6.1)	(3.3)	(4.1)	(100.0)			
合計	1,424	1,112	1,660	2,662	7,124	8,235	4,120	1,850	901	1,239	30,327	311		30,638
(%)	(4.7)	(3.7)	(5.5)	(8.8)	(23.5)	(27.2)	(13.6)	(6.1)	(3.0)	(4.1)	(100.0)			
記載なし												2		2
(%)												(0.0065)		
総計	1,424	1,112	1,660	2,662	7,124	8,235	4,120	1,850	901	1,239	30,327	313		30,640
(%)	(4.7)	(3.7)	(5.5)	(8.8)	(23.5)	(27.2)	(13.6)	(6.1)	(3.0)	(4.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##A0262##
(C)Japanese Society for Dialysis Therapy
2016年調査