

表 328 2016年死亡患者死亡原因の詳細(原疾患との関係)

死因	慢性 糸球体 腎炎	慢性腎盂 腎炎	急速 進行性 糸球体 腎炎	妊婦腎 /妊娠 中毒症	その他の 分類不能 の腎炎	多発性 囊胞腎	腎硬化症	悪性 高血圧	糖尿病 性腎症	SLE腎炎	アミロイド 腎	痛風腎	先天性 代謝異常 による 腎不全	腎・尿路 結核	腎・尿路 結石	腎・尿路 腫瘍	閉塞性 尿路障害	骨髄腫	腎形成 不全	不明	再導入	その他	合計	記載なし	総計
心不全	1,542	53	51	16	20	85	746	34	2,741	48	17	8	3	5	11	13	16	13	5	835	31	201	6,494		6,494
(%)	(23.7)	(0.8)	(0.8)	(0.2)	(0.3)	(1.3)	(11.5)	(0.5)	(42.2)	(0.7)	(0.3)	(0.1)	(0.0)	(0.1)	(0.2)	(0.2)	(0.2)	(0.2)	(0.1)	(12.9)	(0.5)	(3.1)	(100.0)		
肺水腫(溢水)	60	1	4		2	3	32	3	138			1								42	2	13	301		301
(%)	(19.9)	(0.3)	(1.3)		(0.7)	(1.0)	(10.6)	(1.0)	(45.8)			(0.3)								(14.0)	(0.7)	(4.3)	(100.0)		
急性心筋梗塞(発症30日以内死亡)	171	4	4	2	2	19	91	3	414	3		1			1	1	2			87	3	20	828		828
(%)	(20.7)	(0.5)	(0.5)	(0.2)	(0.2)	(2.3)	(11.0)	(0.4)	(50.0)	(0.4)		(0.1)			(0.1)	(0.1)	(0.2)			(10.5)	(0.4)	(2.4)	(100.0)		
虚血性心疾患(急性心筋梗塞以外)	67	2	3			6	43		196	2						1			1	50	3	8	382		382
(%)	(17.5)	(0.5)	(0.8)			(1.6)	(11.3)		(51.3)	(0.5)						(0.3)			(0.3)	(13.1)	(0.8)	(2.1)	(100.0)		
不整脈、伝導障害	78	1	5	1	2	7	48	3	203	2	1	2		1						58	3	16	431		431
(%)	(18.1)	(0.2)	(1.2)	(0.2)	(0.5)	(1.6)	(11.1)	(0.7)	(47.1)	(0.5)	(0.2)	(0.5)		(0.2)						(13.5)	(0.7)	(3.7)	(100.0)		
心内膜炎および弁膜症	51	1		1		4	26	1	60	4	2				2					23	1	10	186		186
(%)	(27.4)	(0.5)		(0.5)		(2.2)	(14.0)	(0.5)	(32.3)	(2.2)	(1.1)				(1.1)					(12.4)	(0.5)	(5.4)	(100.0)		
その他の心疾患	111	5	10	1	4	14	65	6	155	3	5	1	1		1		1		1	68	2	21	475		475
(%)	(23.4)	(1.1)	(2.1)	(0.2)	(0.8)	(2.9)	(13.7)	(1.3)	(32.6)	(0.6)	(1.1)	(0.2)	(0.2)		(0.2)		(0.2)		(0.2)	(14.3)	(0.4)	(4.4)	(100.0)		
くも膜下出血	48		3	1	1	6	19	1	58	1	1		1							14		2	156		156
(%)	(30.8)		(1.9)	(0.6)	(0.6)	(3.8)	(12.2)	(0.6)	(37.2)	(0.6)	(0.6)		(0.6)							(9.0)		(1.3)	(100.0)		
脳内出血	197	6	7	1	1	40	85	7	438	3	1	1		1	2	2	1	1	2	94	3	31	924		924
(%)	(21.3)	(0.6)	(0.8)	(0.1)	(0.1)	(4.3)	(9.2)	(0.8)	(47.4)	(0.3)	(0.1)	(0.1)		(0.1)	(0.2)	(0.2)	(0.1)	(0.1)	(0.2)	(10.2)	(0.3)	(3.4)	(100.0)		
脳梗塞	136	2	10	1	1	12	90	8	320	2	1	1	1					2		102		13	702		702
(%)	(19.4)	(0.3)	(1.4)	(0.1)	(0.1)	(1.7)	(12.8)	(1.1)	(45.6)	(0.3)	(0.1)	(0.1)	(0.1)					(0.3)		(14.5)		(1.9)	(100.0)		
その他の脳血管疾患	51	1	2	1	1	2	33	2	79	1	1	1								18	1	4	197		197
(%)	(25.9)	(0.5)	(1.0)	(0.5)	(0.5)	(1.0)	(16.8)	(1.0)	(40.1)	(0.5)		(0.5)								(9.1)	(0.5)	(2.0)	(100.0)		
敗血症	458	21	32	4	8	79	223	11	1,100	31	9	7	1	2	3	9	3	6		339	19	111	2,476		2,476
(%)	(18.5)	(0.8)	(1.3)	(0.2)	(0.3)	(3.2)	(9.0)	(0.4)	(44.4)	(1.3)	(0.4)	(0.3)	(0.0)	(0.1)	(0.1)	(0.4)	(0.1)	(0.2)		(13.7)	(0.8)	(4.5)	(100.0)		
中枢神経系感染症	9	1	1			2			4	1							1	1		3			23		23
(%)	(39.1)	(4.3)	(4.3)			(8.7)			(17.4)	(4.3)							(4.3)	(4.3)		(13.0)			(100.0)		
肺炎	831	26	69	4	14	60	561	16	1,309	16	7	12	1	5	2	12	7	3	1	496	17	126	3,595		3,595
(%)	(23.1)	(0.7)	(1.9)	(0.1)	(0.4)	(1.7)	(15.6)	(0.4)	(36.4)	(0.4)	(0.2)	(0.3)	(0.0)	(0.1)	(0.1)	(0.3)	(0.2)	(0.1)	(0.0)	(13.8)	(0.5)	(3.5)	(100.0)		
インフルエンザ	7		1		1		1	1	12			1								3		4	31		31
(%)	(22.6)		(3.2)		(3.2)		(3.2)	(3.2)	(38.7)			(3.2)								(9.7)		(12.9)	(100.0)		
尿路感染症	6					2	4		6											6			24		24
(%)	(25.0)					(8.3)	(16.7)		(25.0)											(25.0)			(100.0)		
消化管・胆道系感染症・腹膜炎	56	2	2	1	1	14	33		76	2	1					1		1		30	1	7	228		228
(%)	(24.6)	(0.9)	(0.9)	(0.4)	(0.4)	(6.1)	(14.5)		(33.3)	(0.9)	(0.4)					(0.4)		(0.4)		(13.2)	(0.4)	(3.1)	(100.0)		
劇症(急性)ウイルス肝炎	2					1			1											2		1	8		8
(%)	(25.0)					(12.5)			(12.5)							(12.5)				(25.0)		(12.5)	(100.0)		
結核	6		1			2	9		7	1										13		5	44		44
(%)	(13.6)		(2.3)			(4.5)	(20.5)		(15.9)	(2.3)										(29.5)		(11.4)	(100.0)		
ヒト免疫不全ウイルス[HIV]感染症									1														1		1
(%)									(100.0)														(100.0)		
その他の感染症	56	1	4		1	15	25		115	5		1			1		1	2		58	4	6	295		295
(%)	(19.0)	(0.3)	(1.4)		(0.3)	(5.1)	(8.5)		(39.0)	(1.7)		(0.3)			(0.3)		(0.3)	(0.7)		(19.7)	(1.4)	(2.0)	(100.0)		
中枢神経系の悪性新生物	8	1		1		4			11									1		3	1	1	31		31
(%)	(25.8)	(3.2)		(3.2)		(12.9)			(35.5)									(3.2)		(9.7)	(3.2)	(3.2)	(100.0)		
呼吸器系の悪性新生物	123	3	6			13	74	3	206	4		1			2	6	2			67	2	23	535		535
(%)	(23.0)	(0.6)	(1.1)			(2.4)	(13.8)	(0.6)	(38.5)	(0.7)		(0.2)			(0.4)	(1.1)	(0.4)			(12.5)	(0.4)	(4.3)	(100.0)		
肝癌	85	6	3	1	2	7	36	3	200	1	1	1				2				66		13	428		428
(%)	(19.9)	(1.4)	(0.7)	(0.2)	(0.5)	(1.6)	(8.4)	(0.7)	(46.7)	(0.2)	(0.2)	(0.2)	(0.2)			(0.5)				(15.4)		(3.0)	(100.0)		
肝癌以外の消化器系の悪性新生物	193	7	6	3	3	19	82	3	315	3	2	1		2	3	7	8			106	1	30	794		794
(%)	(24.3)	(0.9)	(0.8)	(0.4)	(0.4)	(2.4)	(10.3)	(0.4)	(39.7)	(0.4)	(0.3)	(0.1)		(0.3)	(0.4)	(0.9)	(1.0)			(13.4)	(0.1)	(3.8)	(100.0)		
乳房の悪性新生物	18					1	3		17	1					1					6	1	3	51		51
(%)	(35.3)					(2.0)	(5.9)		(33.3)	(2.0)					(2.0)					(11.8)	(2.0)	(5.9)	(100.0)		
性器の悪性新生物	19						10	1	22						1	5	1			15		3	77		77
(%)	(24.7)						(13.0)	(1.3)	(28.6)						(1.3)	(6.5)	(1.3)			(19.5)		(3.9)	(100.0)		

表 328 2016年死亡患者死亡原因の詳細(原疾患との関係)

死因	慢性 糸球体 腎炎	慢性腎盂 腎炎	急速 進行性 糸球体 腎炎	妊娠腎 /妊婦 中毒症	その他の 分類不能 の腎炎	多発性 囊胞腎	腎硬化症	悪性 高血圧	糖尿病 性腎症	SLE腎炎	アミロイド 腎	痛風腎	先天性 代謝異常 による 腎不全	腎・尿路 結核	腎・尿路 結石	腎・尿路 腫瘍	閉塞性 尿路障害	骨髄腫	腎形成 不全	不明	再導入	その他	合計	記載なし	総計
腎の悪性新生物 (%)	78 (32.9)	2 (0.8)	1 (0.4)		1 (0.4)	5 (2.1)	29 (12.2)		47 (19.8)			1 (0.4)			2 (0.8)	31 (13.1)	1 (0.4)			22 (9.3)		17 (7.2)	237 (100.0)		237
内分泌腺の悪性新生物 (%)	11 (16.9)					3 (4.6)	14 (21.5)		21 (32.3)	2 (3.1)										12 (18.5)	1 (1.5)	1 (1.5)	65 (100.0)		65
造血・リンパ組織の悪性新生物 (%)	51 (18.6)	8 (2.9)	4 (1.5)	1 (0.4)		4 (1.5)	24 (8.8)	1 (0.4)	64 (23.4)	1 (0.4)	9 (3.3)				1 (0.4)	1 (0.4)	51 (18.6)			39 (14.2)		15 (5.5)	274 (100.0)		274
その他の悪性新生物 (%)	94 (19.8)	9 (1.9)	7 (1.5)		3 (0.6)	14 (2.9)	47 (9.9)	3 (0.6)	153 (32.2)	1 (0.2)	3 (0.6)	5 (1.1)	1 (0.2)	1 (0.2)	2 (0.4)	20 (4.2)	5 (1.1)	8 (1.7)		70 (14.7)	1 (0.2)	28 (5.9)	475 (100.0)		475
ウイルス性肝硬変 (%)	27 (30.7)	2 (2.3)	1 (1.1)		1 (1.1)	2 (2.3)	4 (4.5)		32 (36.4)						2 (2.3)					15 (17.0)		2 (2.3)	88 (100.0)		88
ウイルス性以外の肝硬変 (%)	35 (21.6)	2 (1.2)	3 (1.9)		1 (0.6)	6 (3.7)	16 (9.9)	2 (1.2)	50 (30.9)		1 (0.6)				1 (0.6)					32 (19.8)		13 (8.0)	162 (100.0)		162
膵の血行障害 (%)	46 (24.0)		2 (1.0)		2 (1.0)	3 (1.6)	22 (11.5)	2 (1.0)	75 (39.1)	1 (0.5)							1 (0.5)			30 (15.6)		8 (4.2)	192 (100.0)		192
イレウス (%)	26 (19.4)	1 (0.7)	1 (0.7)	1 (0.7)	2 (1.5)	4 (3.0)	17 (12.7)	1 (0.7)	44 (32.8)	2 (1.5)					1 (0.7)			1 (0.7)	1 (0.7)	23 (17.2)	1 (0.7)	8 (6.0)	134 (100.0)		134
消化管出血 (%)	89 (22.5)	1 (0.3)	8 (2.0)	2 (0.5)		7 (1.8)	53 (13.4)	2 (0.5)	133 (33.7)	2 (0.5)		2 (0.5)	1 (0.3)		2 (0.5)	2 (0.5)	1 (0.3)	2 (0.5)		66 (16.7)	1 (0.3)	21 (5.3)	395 (100.0)		395
被覆性腹膜硬化症 (%)	4 (100.0)																						4 (100.0)		4
その他の消化器疾患 (%)	58 (25.8)	4 (1.8)	2 (0.9)		1 (0.4)	10 (4.4)	30 (13.3)	5 (2.2)	72 (32.0)	1 (0.4)							2 (0.9)	1 (0.4)	1 (0.4)	30 (13.3)	1 (0.4)	7 (3.1)	225 (100.0)		225
肺梗塞、肺塞栓症 (%)	19 (31.7)	1 (1.7)	1 (1.7)				1 (1.7)		18 (30.0)							2 (3.3)		2 (3.3)		11 (18.3)		5 (8.3)	60 (100.0)		60
悪液質 (%)	146 (21.3)	2 (0.3)	11 (1.6)		2 (0.3)	23 (3.4)	96 (14.0)	7 (1.0)	264 (38.6)	7 (1.0)	1 (0.1)	2 (0.3)		1 (0.1)	1 (0.1)	1 (0.1)	1 (0.1)	3 (0.4)		100 (14.6)	2 (0.3)	14 (2.0)	684 (100.0)		684
尿毒症 (%)	74 (19.1)	1 (0.3)	5 (1.3)		2 (0.5)	8 (2.1)	42 (10.8)	4 (1.0)	157 (40.5)	2 (0.5)	2 (0.5)	2 (0.3)	1 (0.3)		1 (0.3)		2 (0.5)	1 (0.3)		78 (20.1)		7 (1.8)	388 (100.0)		388
認知症 (%)	10 (17.2)					1 (1.7)	13 (22.4)		23 (39.7)								1 (1.7)	1 (1.7)		10 (17.2)			58 (100.0)		58
その他の悪液質/尿毒症 (%)	45 (30.2)	1 (0.7)	1 (0.7)	1 (0.7)		1 (0.7)	17 (11.4)		48 (32.2)		2 (1.3)							1 (0.7)	1 (0.7)	27 (18.1)		4 (2.7)	149 (100.0)		149
高カリウム血症 (%)	17 (20.5)	2 (2.4)	1 (1.2)			1 (1.2)	9 (10.8)		36 (43.4)	1 (1.2)								1 (1.2)		9 (10.8)		6 (7.2)	83 (100.0)		83
原因不明の突然死 (%)	144 (21.9)	6 (0.9)	4 (0.6)			8 (1.2)	73 (11.1)	5 (0.8)	330 (50.1)	3 (0.5)	2 (0.3)				2 (0.3)	1 (0.2)	1 (0.2)			68 (10.3)		12 (1.8)	659 (100.0)		659
自殺 (%)	19 (25.7)	2 (2.7)	1 (1.4)		1 (1.4)	3 (4.1)	4 (5.4)		33 (44.6)											9 (12.2)		2 (2.7)	74 (100.0)		74
治療拒否(透析拒否) (%)	30 (24.0)		1 (0.8)		1 (0.8)	3 (2.4)	22 (17.6)	2 (1.6)	41 (32.8)		2 (1.6)						1 (0.8)	1 (0.8)		13 (10.4)		8 (6.4)	125 (100.0)		125
災害・事故死 (%)	44 (27.2)	2 (1.2)		1 (0.6)		4 (2.5)	15 (9.3)		75 (46.3)								2 (1.2)			13 (8.0)		4 (2.5)	162 (100.0)		162
その他 (%)	631 (22.1)	12 (0.4)	25 (0.9)	6 (0.2)	12 (0.4)	62 (2.2)	316 (11.1)	13 (0.5)	1,088 (38.1)	28 (1.0)	10 (0.4)	5 (0.2)	2 (0.1)	2 (0.1)	7 (0.2)	8 (0.3)	5 (0.2)	3 (0.1)	1 (0.0)	429 (15.0)	6 (0.2)	182 (6.4)	2,853 (100.0)		2,853
不明 (%)	696 (20.6)	21 (0.6)	22 (0.7)	11 (0.3)	13 (0.4)	62 (1.8)	350 (10.4)	14 (0.4)	1,433 (42.5)	15 (0.4)	6 (0.2)	11 (0.3)		2 (0.1)	5 (0.1)	11 (0.3)	9 (0.3)	12 (0.4)	3 (0.1)	543 (16.1)	14 (0.4)	122 (3.6)	3,375 (100.0)		3,375
合計 (%)	6,783 (22.1)	223 (0.7)	325 (1.1)	62 (0.2)	106 (0.3)	645 (2.1)	3,559 (11.6)	167 (0.5)	12,441 (40.6)	200 (0.7)	86 (0.3)	67 (0.2)	15 (0.0)	23 (0.1)	57 (0.2)	136 (0.4)	78 (0.3)	116 (0.4)	16 (0.1)	4,253 (13.9)	122 (0.4)	1,158 (3.8)	30,638 (100.0)		30,638
記載なし (%)							2 (100.0)																2 (100.0)		2
総計 (%)	6,783 (22.1)	223 (0.7)	325 (1.1)	62 (0.2)	106 (0.3)	645 (2.1)	3,561 (11.6)	167 (0.5)	12,441 (40.6)	200 (0.7)	86 (0.3)	67 (0.2)	15 (0.0)	23 (0.1)	57 (0.2)	136 (0.4)	78 (0.3)	116 (0.4)	16 (0.1)	4,253 (13.9)	122 (0.4)	1,158 (3.8)	30,640 (100.0)		30,640

数値下のかつこ内は行方向の合計に対する%です。

##A0263##

(C)Japanese Society for Dialysis Therapy

2016年調査