

表 330 1983年以降導入2016年末患者原疾患別人数, 平均年齢

原疾患	人数	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	89,001		89,001	67.33	12.51
(%)	(28.2)		(28.2)		
慢性腎盂腎炎	2,820		2,820	66.20	13.91
(%)	(0.9)		(0.9)		
急速進行性糸球体腎炎	2,521		2,521	69.59	12.72
(%)	(0.8)		(0.8)		
妊娠腎/妊娠中毒症	1,378		1,378	65.50	10.52
(%)	(0.4)		(0.4)		
その他の分類不能の腎炎	1,417		1,417	61.37	17.06
(%)	(0.4)		(0.4)		
多発性嚢胞腎	11,425		11,425	65.41	11.39
(%)	(3.6)		(3.6)		
腎硬化症	31,613		31,613	74.33	11.76
(%)	(10.0)		(10.0)		
悪性高血圧	2,716		2,716	64.21	14.88
(%)	(0.9)		(0.9)		
糖尿病性腎症	123,594		123,594	67.70	11.38
(%)	(39.2)		(39.2)		
SLE腎炎	2,192		2,192	61.76	13.73
(%)	(0.7)		(0.7)		
アミロイド腎	436		436	67.78	11.22
(%)	(0.1)		(0.1)		
痛風腎	1,047		1,047	68.01	11.27
(%)	(0.3)		(0.3)		
先天性代謝異常による腎不全	271		271	50.73	17.55
(%)	(0.1)		(0.1)		
腎・尿路結核	156		156	73.74	8.86
(%)	(0.0)		(0.0)		
腎・尿路結石	572		572	71.24	10.86
(%)	(0.2)		(0.2)		
腎・尿路腫瘍	958		958	72.32	10.78
(%)	(0.3)		(0.3)		
閉塞性尿路障害	736		736	65.02	15.76
(%)	(0.2)		(0.2)		
骨髄腫	323		323	70.95	10.81
(%)	(0.1)		(0.1)		
腎形成不全	642		642	45.60	19.29
(%)	(0.2)		(0.2)		
不明	30,991	2	30,993	70.32	13.11
(%)	(9.8)	(100.0)	(9.8)		
再導入	2,105		2,105	56.88	12.62
(%)	(0.7)		(0.7)		
その他	8,334		8,334	66.20	15.73
(%)	(2.6)		(2.6)		
合計	315,248	2	315,250	68.17	12.52
(%)	(100.0)	(100.0)	(100.0)		
記載なし					
総計	315,248	2	315,250	68.17	12.52

数値下のかっこ内は列方向の合計に対する%です。

##B0002##

(C)Japanese Society for Dialysis Therapy

2016年調査