

表 358 2014年導入患者2年生存率(原疾患別、年齢別)

原疾患	0歳～		15歳～		30歳～		45歳～		60歳～		75歳～		90歳～		記載なし		合計	
慢性糸球体腎炎	2	1.000	59	1.000	388	0.983	1,015	0.965	2,451	0.913	2,927	0.789	253	0.683			7,095	0.865
慢性腎盂腎炎			3	1.000	21	1.000	28	0.890	101	0.900	120	0.617	10	0.500			283	0.771
急速進行性糸球体腎炎	1	1.000	4	1.000	10	0.789	40	0.899	163	0.689	251	0.531	21	0.476			490	0.619
妊娠腎／妊娠中毒症					1	1.000	13	1.000	15	1.000	13	0.923	1	1.000			43	0.976
その他分類不能の腎炎	3	1.000	13	1.000	14	0.929	22	0.909	49	0.899	55	0.653	6	0.666			162	0.820
多発性嚢胞腎			4	1.000	53	0.980	318	0.967	432	0.947	253	0.854	12	0.917			1,072	0.931
腎硬化症			15	0.793	81	0.960	368	0.949	1,371	0.895	3,166	0.775	346	0.679			5,347	0.814
悪性高血圧			2	1.000	47	1.000	71	0.956	93	0.892	115	0.722	11	0.818			339	0.857
糖尿病性腎症	1	1.000	5	0.750	603	0.966	2,935	0.946	7,580	0.886	5,768	0.797	285	0.682			17,177	0.866
SLE腎炎			5	1.000	27	0.872	54	0.883	103	0.793	87	0.579	11	0.546			287	0.743
アミロイド腎					2	0.500	12	0.830	35	0.563	45	0.555	2	1.000			96	0.600
痛風腎					2	1.000	8	1.000	32	0.969	25	0.920	2	1.000			69	0.956
先天性代謝異常に基づく腎不全			3	1.000	1	1.000	2	1.000	6	1.000	5	0.800					17	0.938
腎・尿路結核							2	1.000	2	1.000	5	0.800	2	1.000			11	0.909
腎・尿路結石					5	1.000	7	0.857	29	0.828	31	0.774					72	0.820
腎・尿路腫瘍			1	1.000	2	1.000	9	0.778	71	0.729	100	0.620	7	0.286			190	0.662
閉塞性尿路障害			1	1.000	4	0.750	10	0.684	34	0.736	42	0.735	2	0.000			93	0.715
骨髄腫			2	0.000	2	0.000	8	0.357	62	0.444	58	0.293	3	0.000			135	0.353
腎形成不全	4	1.000	22	1.000	12	1.000	3	1.000	9	0.889	8	0.750	1	0.000			59	0.925
不明	2	0.500	16	0.846	155	0.931	478	0.903	1,433	0.789	2,347	0.678	275	0.594			4,706	0.737
移植後再導入			6	1.000	29	1.000	67	0.922	73	0.800	33	0.576	4	0.500			212	0.826
その他	6	1.000	28	0.915	67	0.868	167	0.781	486	0.674	642	0.551	62	0.536			1,458	0.640
記載なし																		
合計	19	0.946	189	0.937	1,526	0.960	5,637	0.939	14,630	0.870	16,096	0.754	1,316	0.650			39,413	0.829

(C)Japanese Society for Dialysis Therapy

2016年調査