

表 532 糖尿病の既往 原疾患別（腹膜透析患者）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	1,568	302	1,870	13	783	2,666
(%)	(83.9)	(16.1)	(100.0)			
慢性腎盂腎炎	67	8	75		22	97
(%)	(89.3)	(10.7)	(100.0)			
急速進行性糸球体腎炎	34	8	42		19	61
(%)	(81.0)	(19.0)	(100.0)			
妊娠腎/妊娠中毒症	19	2	21		3	24
(%)	(90.5)	(9.5)	(100.0)			
その他の分類不能の腎炎	42	8	50		25	75
(%)	(84.0)	(16.0)	(100.0)			
多発性嚢胞腎	154	26	180	1	91	272
(%)	(85.6)	(14.4)	(100.0)			
腎硬化症	602	166	768	5	333	1,106
(%)	(78.4)	(21.6)	(100.0)			
悪性高血圧	63	10	73		33	106
(%)	(86.3)	(13.7)	(100.0)			
糖尿病性腎症		2,577	2,577		169	2,746
(%)		(100.0)	(100.0)			
SLE腎炎	27	8	35		20	55
(%)	(77.1)	(22.9)	(100.0)			
アミロイド腎	9		9		7	16
(%)	(100.0)		(100.0)			
痛風腎	15	2	17		6	23
(%)	(88.2)	(11.8)	(100.0)			
先天性代謝異常による腎不全	11	2	13		4	17
(%)	(84.6)	(15.4)	(100.0)			
腎・尿路結核					1	1
(%)						
腎・尿路結石	17		17		4	21
(%)	(100.0)		(100.0)			
腎・尿路腫瘍	6	3	9		3	12
(%)	(66.7)	(33.3)	(100.0)			
閉塞性尿路障害	12	1	13		7	20
(%)	(92.3)	(7.7)	(100.0)			
骨髄腫	5	1	6		1	7
(%)	(83.3)	(16.7)	(100.0)			
腎形成不全	43	4	47		18	65
(%)	(91.5)	(8.5)	(100.0)			
不明	431	154	585	7	297	889
(%)	(73.7)	(26.3)	(100.0)			
再導入	42	8	50		15	65
(%)	(84.0)	(16.0)	(100.0)			
その他	139	40	179	1	91	271
(%)	(77.7)	(22.3)	(100.0)			
合計	3,306	3,330	6,636	27	1,952	8,615
(%)	(49.8)	(50.2)	(100.0)			
記載なし						
(%)						
総計	3,306	3,330	6,636	27	1,952	8,615
(%)	(49.8)	(50.2)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D0043##

(C)Japanese Society for Dialysis Therapy

2016年調査