

表 540 心筋梗塞の既往 原疾患別 (透析患者全体)

| 原疾患           | 既往なし    | 既往あり   | 合計      | 不明    | 記載なし   | 総計      |
|---------------|---------|--------|---------|-------|--------|---------|
| 慢性糸球体腎炎       | 63,128  | 5,214  | 68,342  | 272   | 23,169 | 91,783  |
| (%)           | (92.4)  | (7.6)  | (100.0) |       |        |         |
| 慢性腎盂腎炎        | 2,051   | 149    | 2,200   | 7     | 689    | 2,896   |
| (%)           | (93.2)  | (6.8)  | (100.0) |       |        |         |
| 急速進行性糸球体腎炎    | 1,771   | 132    | 1,903   | 9     | 626    | 2,538   |
| (%)           | (93.1)  | (6.9)  | (100.0) |       |        |         |
| 妊娠腎/妊娠中毒症     | 1,080   | 64     | 1,144   | 6     | 333    | 1,483   |
| (%)           | (94.4)  | (5.6)  | (100.0) |       |        |         |
| その他の分類不能の腎炎   | 971     | 68     | 1,039   | 5     | 410    | 1,454   |
| (%)           | (93.5)  | (6.5)  | (100.0) |       |        |         |
| 多発性嚢胞腎        | 8,209   | 539    | 8,748   | 43    | 2,680  | 11,471  |
| (%)           | (93.8)  | (6.2)  | (100.0) |       |        |         |
| 腎硬化症          | 21,578  | 2,466  | 24,044  | 113   | 7,493  | 31,650  |
| (%)           | (89.7)  | (10.3) | (100.0) |       |        |         |
| 悪性高血圧         | 1,853   | 172    | 2,025   | 25    | 671    | 2,721   |
| (%)           | (91.5)  | (8.5)  | (100.0) |       |        |         |
| 糖尿病性腎症        | 79,562  | 13,293 | 92,855  | 460   | 30,342 | 123,657 |
| (%)           | (85.7)  | (14.3) | (100.0) |       |        |         |
| SLE腎炎         | 1,565   | 104    | 1,669   | 7     | 546    | 2,222   |
| (%)           | (93.8)  | (6.2)  | (100.0) |       |        |         |
| アミロイド腎        | 306     | 20     | 326     | 2     | 110    | 438     |
| (%)           | (93.9)  | (6.1)  | (100.0) |       |        |         |
| 痛風腎           | 767     | 61     | 828     | 5     | 223    | 1,056   |
| (%)           | (92.6)  | (7.4)  | (100.0) |       |        |         |
| 先天性代謝異常による腎不全 | 208     | 10     | 218     |       | 63     | 281     |
| (%)           | (95.4)  | (4.6)  | (100.0) |       |        |         |
| 腎・尿路結核        | 113     | 7      | 120     |       | 48     | 168     |
| (%)           | (94.2)  | (5.8)  | (100.0) |       |        |         |
| 腎・尿路結石        | 423     | 30     | 453     | 4     | 121    | 578     |
| (%)           | (93.4)  | (6.6)  | (100.0) |       |        |         |
| 腎・尿路腫瘍        | 654     | 60     | 714     | 4     | 241    | 959     |
| (%)           | (91.6)  | (8.4)  | (100.0) |       |        |         |
| 閉塞性尿路障害       | 537     | 25     | 562     | 1     | 180    | 743     |
| (%)           | (95.6)  | (4.4)  | (100.0) |       |        |         |
| 骨髄腫           | 233     | 14     | 247     | 2     | 74     | 323     |
| (%)           | (94.3)  | (5.7)  | (100.0) |       |        |         |
| 腎形成不全         | 492     | 12     | 504     |       | 160    | 664     |
| (%)           | (97.6)  | (2.4)  | (100.0) |       |        |         |
| 不明            | 20,593  | 1,975  | 22,568  | 186   | 8,658  | 31,412  |
| (%)           | (91.2)  | (8.8)  | (100.0) |       |        |         |
| 再導入           | 1,529   | 112    | 1,641   | 6     | 527    | 2,174   |
| (%)           | (93.2)  | (6.8)  | (100.0) |       |        |         |
| その他           | 5,611   | 519    | 6,130   | 38    | 2,270  | 8,438   |
| (%)           | (91.5)  | (8.5)  | (100.0) |       |        |         |
| 合計            | 213,234 | 25,046 | 238,280 | 1,195 | 79,634 | 319,109 |
| (%)           | (89.5)  | (10.5) | (100.0) |       |        |         |
| 記載なし          |         |        |         |       |        |         |
| (%)           |         |        |         |       |        |         |
| 総計            | 213,234 | 25,046 | 238,280 | 1,195 | 79,634 | 319,109 |
| (%)           | (89.5)  | (10.5) | (100.0) |       |        |         |

数値下のかっこ内は行方向の合計に対する%です。

##D0116##

(C)Japanese Society for Dialysis Therapy

2016年調査