

表 550 心筋梗塞の既往 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	16,669	1,416	18,085	58	4,245	22,388
(%)	(92.2)	(7.8)	(100.0)			
慢性腎盂腎炎	472	35	507		120	627
(%)	(93.1)	(6.9)	(100.0)			
急速進行性糸球体腎炎	362	24	386	2	96	484
(%)	(93.8)	(6.2)	(100.0)			
妊娠腎/妊娠中毒症	349	23	372		87	459
(%)	(93.8)	(6.2)	(100.0)			
その他の分類不能の腎炎	264	16	280		75	355
(%)	(94.3)	(5.7)	(100.0)			
多発性嚢胞腎	1,976	138	2,114	10	445	2,569
(%)	(93.5)	(6.5)	(100.0)			
腎硬化症	4,346	498	4,844	18	1,034	5,896
(%)	(89.7)	(10.3)	(100.0)			
悪性高血圧	408	42	450	2	103	555
(%)	(90.7)	(9.3)	(100.0)			
糖尿病性腎症	17,087	2,808	19,895	75	4,546	24,516
(%)	(85.9)	(14.1)	(100.0)			
SLE腎炎	417	29	446	2	89	537
(%)	(93.5)	(6.5)	(100.0)			
アミロイド腎	73	7	80	1	17	98
(%)	(91.3)	(8.8)	(100.0)			
痛風腎	166	19	185	1	41	227
(%)	(89.7)	(10.3)	(100.0)			
先天性代謝異常による腎不全	58	3	61		12	73
(%)	(95.1)	(4.9)	(100.0)			
腎・尿路結核	37	1	38		6	44
(%)	(97.4)	(2.6)	(100.0)			
腎・尿路結石	77	10	87		26	113
(%)	(88.5)	(11.5)	(100.0)			
腎・尿路腫瘍	143	14	157		33	190
(%)	(91.1)	(8.9)	(100.0)			
閉塞性尿路障害	115	10	125		26	151
(%)	(92.0)	(8.0)	(100.0)			
骨髄腫	28	2	30		7	37
(%)	(93.3)	(6.7)	(100.0)			
腎形成不全	140	3	143		28	171
(%)	(97.9)	(2.1)	(100.0)			
不明	4,712	427	5,139	11	1,381	6,531
(%)	(91.7)	(8.3)	(100.0)			
再導入	506	31	537	1	95	633
(%)	(94.2)	(5.8)	(100.0)			
その他	1,259	133	1,392	2	326	1,720
(%)	(90.4)	(9.6)	(100.0)			
合計	49,664	5,689	55,353	183	12,838	68,374
(%)	(89.7)	(10.3)	(100.0)			
記載なし						
(%)						
総計	49,664	5,689	55,353	183	12,838	68,374
(%)	(89.7)	(10.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D0133##

(C)Japanese Society for Dialysis Therapy

2016年調査