

表 573 脳出血の既往 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	16,960	1,016	17,976	56	4,356	22,388
(%)	(94.3)	(5.7)	(100.0)			
慢性腎盂腎炎	481	26	507		120	627
(%)	(94.9)	(5.1)	(100.0)			
急速進行性糸球体腎炎	364	19	383	1	100	484
(%)	(95.0)	(5.0)	(100.0)			
妊娠腎/妊娠中毒症	344	26	370		89	459
(%)	(93.0)	(7.0)	(100.0)			
その他の分類不能の腎炎	267	15	282		73	355
(%)	(94.7)	(5.3)	(100.0)			
多発性嚢胞腎	1,892	236	2,128	5	436	2,569
(%)	(88.9)	(11.1)	(100.0)			
腎硬化症	4,460	333	4,793	24	1,079	5,896
(%)	(93.1)	(6.9)	(100.0)			
悪性高血圧	407	46	453	1	101	555
(%)	(89.8)	(10.2)	(100.0)			
糖尿病性腎症	18,523	1,083	19,606	82	4,828	24,516
(%)	(94.5)	(5.5)	(100.0)			
SLE腎炎	431	14	445	2	90	537
(%)	(96.9)	(3.1)	(100.0)			
アミロイド腎	73	4	77	1	20	98
(%)	(94.8)	(5.2)	(100.0)			
痛風腎	175	9	184	1	42	227
(%)	(95.1)	(4.9)	(100.0)			
先天性代謝異常による腎不全	59	2	61		12	73
(%)	(96.7)	(3.3)	(100.0)			
腎・尿路結核	35	3	38		6	44
(%)	(92.1)	(7.9)	(100.0)			
腎・尿路結石	84	4	88		25	113
(%)	(95.5)	(4.5)	(100.0)			
腎・尿路腫瘍	151	6	157		33	190
(%)	(96.2)	(3.8)	(100.0)			
閉塞性尿路障害	115	7	122		29	151
(%)	(94.3)	(5.7)	(100.0)			
骨髄腫	29		29		8	37
(%)	(100.0)		(100.0)			
腎形成不全	139	4	143		28	171
(%)	(97.2)	(2.8)	(100.0)			
不明	4,804	315	5,119	11	1,401	6,531
(%)	(93.8)	(6.2)	(100.0)			
再導入	510	25	535	1	97	633
(%)	(95.3)	(4.7)	(100.0)			
その他	1,263	102	1,365	4	351	1,720
(%)	(92.5)	(7.5)	(100.0)			
合計	51,566	3,295	54,861	189	13,324	68,374
(%)	(94.0)	(6.0)	(100.0)			
記載なし						
(%)						
総計	51,566	3,295	54,861	189	13,324	68,374
(%)	(94.0)	(6.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D0233##

(C)Japanese Society for Dialysis Therapy

2016年調査