

表 586 脳梗塞の既往 原疾患別 (透析患者全体)

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	58,623	10,431	69,054	244	22,485	91,783
(%)	(84.9)	(15.1)	(100.0)			
慢性腎盂腎炎	1,904	315	2,219	11	666	2,896
(%)	(85.8)	(14.2)	(100.0)			
急速進行性糸球体腎炎	1,670	249	1,919	7	612	2,538
(%)	(87.0)	(13.0)	(100.0)			
妊娠腎/妊娠中毒症	982	171	1,153	6	324	1,483
(%)	(85.2)	(14.8)	(100.0)			
その他の分類不能の腎炎	928	123	1,051	4	399	1,454
(%)	(88.3)	(11.7)	(100.0)			
多発性嚢胞腎	7,573	1,269	8,842	36	2,593	11,471
(%)	(85.6)	(14.4)	(100.0)			
腎硬化症	19,482	4,837	24,319	105	7,226	31,650
(%)	(80.1)	(19.9)	(100.0)			
悪性高血圧	1,716	343	2,059	24	638	2,721
(%)	(83.3)	(16.7)	(100.0)			
糖尿病性腎症	72,609	21,316	93,925	402	29,330	123,657
(%)	(77.3)	(22.7)	(100.0)			
SLE腎炎	1,454	229	1,683	8	531	2,222
(%)	(86.4)	(13.6)	(100.0)			
アミロイド腎	294	31	325	2	111	438
(%)	(90.5)	(9.5)	(100.0)			
痛風腎	719	114	833	5	218	1,056
(%)	(86.3)	(13.7)	(100.0)			
先天性代謝異常による腎不全	190	26	216		65	281
(%)	(88.0)	(12.0)	(100.0)			
腎・尿路結核	104	18	122		46	168
(%)	(85.2)	(14.8)	(100.0)			
腎・尿路結石	387	73	460	4	114	578
(%)	(84.1)	(15.9)	(100.0)			
腎・尿路腫瘍	640	88	728	4	227	959
(%)	(87.9)	(12.1)	(100.0)			
閉塞性尿路障害	506	58	564		179	743
(%)	(89.7)	(10.3)	(100.0)			
骨髄腫	229	17	246	2	75	323
(%)	(93.1)	(6.9)	(100.0)			
腎形成不全	479	26	505	1	158	664
(%)	(94.9)	(5.1)	(100.0)			
不明	18,928	3,992	22,920	128	8,364	31,412
(%)	(82.6)	(17.4)	(100.0)			
再導入	1,475	168	1,643	6	525	2,174
(%)	(89.8)	(10.2)	(100.0)			
その他	5,256	919	6,175	35	2,228	8,438
(%)	(85.1)	(14.9)	(100.0)			
合計	196,148	44,813	240,961	1,034	77,114	319,109
(%)	(81.4)	(18.6)	(100.0)			
記載なし						
(%)						
総計	196,148	44,813	240,961	1,034	77,114	319,109
(%)	(81.4)	(18.6)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D0316##

(C)Japanese Society for Dialysis Therapy

2016年調査