

表 591 脳梗塞の既往 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	38,989	7,059	46,048	158	10,966	57,172
(%)	(84.7)	(15.3)	(100.0)			
慢性腎孟腎炎	1,300	214	1,514	9	345	1,868
(%)	(85.9)	(14.1)	(100.0)			
急速進行性糸球体腎炎	1,207	180	1,387	6	294	1,687
(%)	(87.0)	(13.0)	(100.0)			
妊娠腎/妊娠中毒症	595	95	690	6	153	849
(%)	(86.2)	(13.8)	(100.0)			
その他の分類不能の腎炎	595	88	683	3	216	902
(%)	(87.1)	(12.9)	(100.0)			
多発性囊胞腎	5,250	896	6,146	23	1,379	7,548
(%)	(85.4)	(14.6)	(100.0)			
腎硬化症	14,158	3,525	17,683	70	3,765	21,518
(%)	(80.1)	(19.9)	(100.0)			
悪性高血圧	1,196	256	1,452	12	313	1,777
(%)	(82.4)	(17.6)	(100.0)			
糖尿病性腎症	53,155	15,592	68,747	272	15,449	84,468
(%)	(77.3)	(22.7)	(100.0)			
SLE腎炎	948	162	1,110	3	257	1,370
(%)	(85.4)	(14.6)	(100.0)			
アミロイド腎	208	22	230	1	49	280
(%)	(90.4)	(9.6)	(100.0)			
痛風腎	520	82	602	2	96	700
(%)	(86.4)	(13.6)	(100.0)			
先天性代謝異常による腎不全	118	18	136		33	169
(%)	(86.8)	(13.2)	(100.0)			
腎・尿路結核	71	10	81		25	106
(%)	(87.7)	(12.3)	(100.0)			
腎・尿路結石	279	49	328	3	53	384
(%)	(85.1)	(14.9)	(100.0)			
腎・尿路腫瘍	469	65	534	4	106	644
(%)	(87.8)	(12.2)	(100.0)			
閉塞性尿路障害	345	40	385		80	465
(%)	(89.6)	(10.4)	(100.0)			
骨髓腫	167	14	181	1	31	213
(%)	(92.3)	(7.7)	(100.0)			
腎形成不全	283	17	300	1	67	368
(%)	(94.3)	(5.7)	(100.0)			
不明	13,279	2,850	16,129	99	4,250	20,478
(%)	(82.3)	(17.7)	(100.0)			
再導入	852	105	957	4	203	1,164
(%)	(89.0)	(11.0)	(100.0)			
その他	3,633	622	4,255	31	1,120	5,406
(%)	(85.4)	(14.6)	(100.0)			
合計	137,617	31,961	169,578	708	39,250	209,536
(%)	(81.2)	(18.8)	(100.0)			
記載なし						
(%)						
総計	137,617	31,961	169,578	708	39,250	209,536
(%)	(81.2)	(18.8)	(100.0)			

数値下の括弧内は行方向の合計に対する%です。

##D0323##

(C)Japanese Society for Dialysis Therapy

2016年調査