

表 609 四肢切断の有無 原疾患別 (透析患者全体)

原疾患	切断なし	切断あり	合計	不明	記載なし	総計
慢性糸球体腎炎	67,442	1,000	68,442	90	23,251	91,783
(%)	(98.5)	(1.5)	(100.0)			
慢性腎盂腎炎	2,163	43	2,206	2	688	2,896
(%)	(98.1)	(1.9)	(100.0)			
急速進行性糸球体腎炎	1,887	22	1,909	3	626	2,538
(%)	(98.8)	(1.2)	(100.0)			
妊娠腎/妊娠中毒症	1,114	22	1,136	4	343	1,483
(%)	(98.1)	(1.9)	(100.0)			
その他の分類不能の腎炎	1,029	8	1,037	4	413	1,454
(%)	(99.2)	(0.8)	(100.0)			
多発性嚢胞腎	8,676	80	8,756	15	2,700	11,471
(%)	(99.1)	(0.9)	(100.0)			
腎硬化症	23,574	325	23,899	39	7,712	31,650
(%)	(98.6)	(1.4)	(100.0)			
悪性高血圧	2,001	28	2,029	21	671	2,721
(%)	(98.6)	(1.4)	(100.0)			
糖尿病性腎症	85,741	7,011	92,752	166	30,739	123,657
(%)	(92.4)	(7.6)	(100.0)			
SLE腎炎	1,630	40	1,670	3	549	2,222
(%)	(97.6)	(2.4)	(100.0)			
アミロイド腎	323	1	324	1	113	438
(%)	(99.7)	(0.3)	(100.0)			
痛風腎	816	13	829		227	1,056
(%)	(98.4)	(1.6)	(100.0)			
先天性代謝異常による腎不全	215	4	219		62	281
(%)	(98.2)	(1.8)	(100.0)			
腎・尿路結核	119	1	120		48	168
(%)	(99.2)	(0.8)	(100.0)			
腎・尿路結石	451	7	458		120	578
(%)	(98.5)	(1.5)	(100.0)			
腎・尿路腫瘍	704	12	716	2	241	959
(%)	(98.3)	(1.7)	(100.0)			
閉塞性尿路障害	544	16	560		183	743
(%)	(97.1)	(2.9)	(100.0)			
骨髄腫	247		247	2	74	323
(%)	(100.0)		(100.0)			
腎形成不全	499	6	505		159	664
(%)	(98.8)	(1.2)	(100.0)			
不明	22,191	491	22,682	28	8,702	31,412
(%)	(97.8)	(2.2)	(100.0)			
再導入	1,606	46	1,652	1	521	2,174
(%)	(97.2)	(2.8)	(100.0)			
その他	5,980	150	6,130	17	2,291	8,438
(%)	(97.6)	(2.4)	(100.0)			
合計	228,952	9,326	238,278	398	80,433	319,109
(%)	(96.1)	(3.9)	(100.0)			
記載なし						
(%)						
総計	228,952	9,326	238,278	398	80,433	319,109
(%)	(96.1)	(3.9)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D0416##

(C)Japanese Society for Dialysis Therapy

2016年調査