

表 728 腎移植の回数 原疾患別 (透析患者全体)

原疾患	なし	1回	2回	3回	4回以上	合計	不明	記載なし	総計
慢性糸球体腎炎	65,445	924	59	1		66,429	76	25,278	91,783
(%)	(98.5)	(1.4)	(0.1)	(0.0)		(100.0)			
慢性腎盂腎炎	2,130	22	2			2,154	4	738	2,896
(%)	(98.9)	(1.0)	(0.1)			(100.0)			
急速進行性糸球体腎炎	1,816	11				1,827	1	710	2,538
(%)	(99.4)	(0.6)				(100.0)			
妊娠腎/妊娠中毒症	1,049	15	1			1,065	2	416	1,483
(%)	(98.5)	(1.4)	(0.1)			(100.0)			
その他の分類不能の腎炎	998	28	1			1,027	5	422	1,454
(%)	(97.2)	(2.7)	(0.1)			(100.0)			
多発性嚢胞腎	8,323	31				8,354	19	3,098	11,471
(%)	(99.6)	(0.4)				(100.0)			
腎硬化症	22,842	32	2			22,876	54	8,720	31,650
(%)	(99.9)	(0.1)	(0.0)			(100.0)			
悪性高血圧	1,956	6				1,962	4	755	2,721
(%)	(99.7)	(0.3)				(100.0)			
糖尿病性腎症	88,758	146	13			88,917	148	34,592	123,657
(%)	(99.8)	(0.2)	(0.0)			(100.0)			
SLE腎炎	1,607	18				1,625	5	592	2,222
(%)	(98.9)	(1.1)				(100.0)			
アミロイド腎	318					318	1	119	438
(%)	(100.0)					(100.0)			
痛風腎	791	3	1			795	3	258	1,056
(%)	(99.5)	(0.4)	(0.1)			(100.0)			
先天性代謝異常による腎不全	200	10				210		71	281
(%)	(95.2)	(4.8)				(100.0)			
腎・尿路結核	117	1				118		50	168
(%)	(99.2)	(0.8)				(100.0)			
腎・尿路結石	434	1				435	1	142	578
(%)	(99.8)	(0.2)				(100.0)			
腎・尿路腫瘍	711	2				713	1	245	959
(%)	(99.7)	(0.3)				(100.0)			
閉塞性尿路障害	532	8	1			541	1	201	743
(%)	(98.3)	(1.5)	(0.2)			(100.0)			
骨髄腫	237	1				238		85	323
(%)	(99.6)	(0.4)				(100.0)			
腎形成不全	445	37	2			484		180	664
(%)	(91.9)	(7.6)	(0.4)			(100.0)			
不明	21,541	172	16	1		21,730	94	9,588	31,412
(%)	(99.1)	(0.8)	(0.1)	(0.0)		(100.0)			
再導入	552	1,102	53	2		1,709	2	463	2,174
(%)	(32.3)	(64.5)	(3.1)	(0.1)		(100.0)			
その他	5,759	99	8			5,866	7	2,565	8,438
(%)	(98.2)	(1.7)	(0.1)			(100.0)			
合計	226,561	2,669	159	4		229,393	428	89,288	319,109
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			
記載なし									
(%)									
総計	226,561	2,669	159	4		229,393	428	89,288	319,109
(%)	(98.8)	(1.2)	(0.1)	(0.0)		(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D1016##

(C)Japanese Society for Dialysis Therapy

2016年調査