

表 770 透析時間(分/回) 透析歴(年)別 (血液透析濾過患者 透析回数週3回)

| 透析歴 | 180未満 | 180～ | 210～ | 240～ | 270～ | 300～ | 330～ | 360～ | 390～ | 420～ | 合計 | 記載なし | 総計 | 平均 | 標準偏差 |
|------|-------|--------|--------|--------|--------|--------|-------|-------|-------|-------|---------|-------|--------|--------|-------|
| 2年未満 | 32 | 1,526 | 1,028 | 6,567 | 386 | 385 | 15 | 25 | 2 | 27 | 9,993 | 25 | 10,018 | 233.30 | 33.95 |
| (%) | (0.3) | (15.3) | (10.3) | (65.7) | (3.9) | (3.9) | (0.2) | (0.3) | (0.0) | (0.3) | (100.0) | | | | |
| 2年～ | 31 | 1,334 | 1,399 | 11,114 | 950 | 939 | 38 | 79 | 2 | 24 | 15,910 | 28 | 15,938 | 239.94 | 30.45 |
| (%) | (0.2) | (8.4) | (8.8) | (69.9) | (6.0) | (5.9) | (0.2) | (0.5) | (0.0) | (0.2) | (100.0) | | | | |
| 5年～ | 27 | 911 | 1,254 | 12,337 | 1,348 | 1,491 | 50 | 95 | 2 | 27 | 17,542 | 40 | 17,582 | 244.43 | 29.61 |
| (%) | (0.2) | (5.2) | (7.1) | (70.3) | (7.7) | (8.5) | (0.3) | (0.5) | (0.0) | (0.2) | (100.0) | | | | |
| 10年～ | 11 | 279 | 572 | 6,989 | 1,051 | 1,065 | 49 | 60 | 3 | 13 | 10,092 | 24 | 10,116 | 248.52 | 28.00 |
| (%) | (0.1) | (2.8) | (5.7) | (69.3) | (10.4) | (10.6) | (0.5) | (0.6) | (0.0) | (0.1) | (100.0) | | | | |
| 15年～ | | 128 | 243 | 4,115 | 705 | 756 | 41 | 49 | 2 | 5 | 6,044 | 14 | 6,058 | 251.44 | 27.80 |
| (%) | | (2.1) | (4.0) | (68.1) | (11.7) | (12.5) | (0.7) | (0.8) | (0.0) | (0.1) | (100.0) | | | | |
| 20年～ | 1 | 57 | 116 | 2,604 | 493 | 498 | 29 | 36 | 5 | 3 | 3,842 | 15 | 3,857 | 253.07 | 27.95 |
| (%) | (0.0) | (1.5) | (3.0) | (67.8) | (12.8) | (13.0) | (0.8) | (0.9) | (0.1) | (0.1) | (100.0) | | | | |
| 25年～ | 3 | 34 | 64 | 1,503 | 328 | 349 | 24 | 30 | 2 | 1 | 2,338 | 7 | 2,345 | 255.17 | 29.12 |
| (%) | (0.1) | (1.5) | (2.7) | (64.3) | (14.0) | (14.9) | (1.0) | (1.3) | (0.1) | (0.0) | (100.0) | | | | |
| 30年～ | 1 | 25 | 47 | 833 | 201 | 229 | 17 | 10 | | 1 | 1,364 | 3 | 1,367 | 255.72 | 29.03 |
| (%) | (0.1) | (1.8) | (3.4) | (61.1) | (14.7) | (16.8) | (1.2) | (0.7) | | (0.1) | (100.0) | | | | |
| 35年～ | | 13 | 26 | 480 | 112 | 120 | 8 | 18 | 1 | | 778 | 2 | 780 | 257.09 | 31.15 |
| (%) | | (1.7) | (3.3) | (61.7) | (14.4) | (15.4) | (1.0) | (2.3) | (0.1) | | (100.0) | | | | |
| 40年～ | | 2 | 14 | 179 | 47 | 38 | 4 | 2 | | | 286 | 1 | 287 | 254.44 | 27.27 |
| (%) | | (0.7) | (4.9) | (62.6) | (16.4) | (13.3) | (1.4) | (0.7) | | | (100.0) | | | | |
| 合計 | 106 | 4,309 | 4,763 | 46,721 | 5,621 | 5,870 | 275 | 404 | 19 | 101 | 68,189 | 159 | 68,348 | 244.24 | 30.69 |
| (%) | (0.2) | (6.3) | (7.0) | (68.5) | (8.2) | (8.6) | (0.4) | (0.6) | (0.0) | (0.1) | (100.0) | | | | |
| 不明 | | 2 | 1 | 18 | 1 | 2 | | | | | 24 | 2 | 26 | 242.33 | 26.81 |
| (%) | | (8.3) | (4.2) | (75.0) | (4.2) | (8.3) | | | | | (100.0) | | | | |
| 記載なし | | | | | | | | | | | | | | | |
| (%) | | | | | | | | | | | | | | | |
| 総計 | 106 | 4,311 | 4,764 | 46,739 | 5,622 | 5,872 | 275 | 404 | 19 | 101 | 68,213 | 161 | 68,374 | 244.24 | 30.69 |
| (%) | (0.2) | (6.3) | (7.0) | (68.5) | (8.2) | (8.6) | (0.4) | (0.6) | (0.0) | (0.1) | (100.0) | | | | |
| 平均 | 5.07 | 4.51 | 6.52 | 9.01 | 12.25 | 12.28 | 14.70 | 12.66 | 15.89 | 6.16 | 9.13 | 10.16 | 9.14 | | |
| 標準偏差 | 6.09 | 5.70 | 6.86 | 8.26 | 9.37 | 9.34 | 10.42 | 10.04 | 10.46 | 6.23 | 8.47 | 9.00 | 8.47 | | |

数値下のかっこ内は行方向の合計に対する%です。

##D1222##

(C)Japanese Society for Dialysis Therapy

2016年調査