

表 771 透析時間(分/回) 原疾患別 (血液透析濾過患者 透析回数週3回)

原疾患	180未満	180~	210~	240~	270~	300~	330~	360~	390~	420~	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎 (%)	25 (0.1)	1,012 (4.5)	1,269 (5.7)	14,896 (66.7)	2,294 (10.3)	2,487 (11.1)	141 (0.6)	170 (0.8)	12 (0.1)	26 (0.1)	22,332 (100.0)	56	22,388	248.23	30.37
慢性腎盂腎炎 (%)	41 (6.6)	30 (4.8)	451 (72.3)	44 (7.1)	50 (8.0)	5 (0.8)	3 (0.5)				624 (100.0)	3	627	243.69	27.75
急速進行性糸球体腎炎 (%)	39 (8.1)	30 (6.2)	345 (71.3)	36 (7.4)	29 (6.0)	1 (0.2)	4 (0.8)				484 (100.0)	484	484	241.31	27.72
妊娠腎/妊娠中毒症 (%)	1 (0.2)	11 (2.4)	25 (5.4)	318 (69.3)	59 (12.9)	41 (8.9)	3 (0.7)	1 (0.2)			459 (100.0)	459	459	247.88	25.53
その他の分類不能の腎炎 (%)	5 (1.4)	15 (4.2)	27 (7.6)	225 (63.7)	34 (9.6)	44 (12.5)	2 (0.6)	1 (0.3)			353 (100.0)	2	355	247.31	29.46
多発性囊胞腎 (%)	1 (0.0)	146 (5.7)	178 (6.9)	1,783 (69.6)	248 (9.7)	180 (7.0)	4 (0.2)	21 (0.8)	2 (0.1)		2,563 (100.0)	6	2,569	243.97	27.41
腎硬化症 (%)	18 (0.3)	509 (8.6)	542 (9.2)	4,101 (69.7)	349 (5.9)	329 (5.6)	13 (0.2)	18 (0.3)	7 (0.1)		5,886 (100.0)	10	5,896	238.84	29.18
悪性高血圧 (%)	1 (0.2)	32 (5.8)	49 (8.8)	377 (68.1)	42 (7.6)	49 (8.8)	2 (0.4)	1 (0.2)	1 (0.2)		554 (100.0)	1	555	243.02	27.86
糖尿病性腎症 (%)	35 (0.1)	1,666 (6.8)	1,795 (7.3)	17,112 (70.0)	1,743 (7.1)	1,881 (7.7)	66 (0.3)	104 (0.4)	3 (0.0)		24,453 (100.0)	63	24,516	242.79	31.12
SLE腎炎 (%)	1 (0.2)	29 (5.4)	40 (7.4)	363 (67.6)	46 (8.6)	49 (9.1)	2 (0.4)	6 (1.1)			537 (100.0)	537	537	245.83	32.95
アミロイド腎 (%)	11 (11.3)	11 (11.3)	59 (60.8)	9 (9.3)	7 (7.2)						97 (100.0)	1	98	237.78	29.09
痛風腎 (%)	1 (0.4)	11 (4.9)	10 (4.4)	163 (72.4)	16 (7.1)	21 (9.3)	1 (0.4)	2 (0.9)			225 (100.0)	2	227	245.15	27.81
先天性代謝異常による腎不全 (%)	4 (5.5)	6 (8.2)	47 (64.4)	5 (6.8)	10 (13.7)	1 (1.4)					73 (100.0)	73	73	246.42	29.88
腎・尿路結核 (%)	2 (4.5)	5 (11.4)	27 (61.4)	5 (11.4)	5 (11.4)						44 (100.0)	44	44	244.66	28.37
腎・尿路結石 (%)	8 (7.1)	10 (8.9)	81 (72.3)	10 (8.9)	3 (2.7)						112 (100.0)	1	113	238.14	22.63
腎・尿路腫瘍 (%)	11 (5.9)	13 (6.9)	141 (75.0)	7 (3.7)	14 (7.4)						188 (100.0)	2	190	241.73	27.06
閉塞性尿路障害 (%)	15 (9.9)	11 (7.3)	102 (67.5)	10 (6.6)	13 (8.6)						151 (100.0)	151	151	240.38	28.06
骨髄腫 (%)	7 (18.9)	3 (8.1)	24 (64.9)	2 (5.4)	1 (2.7)						37 (100.0)	37	37	233.84	34.20
腎形成不全 (%)	4 (2.3)	9 (5.3)	113 (66.1)	21 (12.3)	23 (13.5)						171 (100.0)	171	171	251.22	27.15
不明 (%)	15 (0.2)	588 (9.0)	545 (8.4)	4,425 (67.8)	440 (6.7)	420 (6.4)	25 (0.4)	47 (0.7)	2 (0.0)		6,522 (100.0)	9	6,531	240.62	31.98
再導入 (%)	1 (0.2)	18 (2.9)	35 (5.6)	407 (64.7)	76 (12.1)	80 (12.7)	2 (0.3)	10 (1.6)			629 (100.0)	4	633	250.88	29.96
その他 (%)	2 (0.1)	132 (7.7)	121 (7.0)	1,179 (68.6)	128 (7.4)	135 (7.9)	6 (0.3)	13 (0.8)	3 (0.2)		1,719 (100.0)	1	1,720	243.35	32.27
合計 (%)	106 (0.2)	4,311 (6.3)	4,764 (7.0)	46,739 (68.5)	5,622 (8.2)	5,872 (8.6)	275 (0.4)	404 (0.6)	19 (0.0)		68,213 (100.0)	161	68,374	244.24	30.69
記載なし (%)															
総計 (%)	106 (0.2)	4,311 (6.3)	4,764 (7.0)	46,739 (68.5)	5,622 (8.2)	5,872 (8.6)	275 (0.4)	404 (0.6)	19 (0.0)		68,213 (100.0)	161	68,374	244.24	30.69

数値下のかっこ内は行方向の合計に対する%です。

##D1223##

(C)Japanese Society for Dialysis Therapy

2016年調査