

表 812 HDF希釈方法 透析前リン濃度(mg/dL)別 (オフラインHDF 透析回数週3回)

透析前リン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3未満	13	148			161		22	183
(%)	(8.1)	(91.9)			(100.0)			
3.0～	21	186			207		31	238
(%)	(10.1)	(89.9)			(100.0)			
3.5～	54	303			357		32	389
(%)	(15.1)	(84.9)			(100.0)			
4.0～	64	393		1	458		50	508
(%)	(14.0)	(85.8)		(0.2)	(100.0)			
4.5～	78	448	1		527		65	592
(%)	(14.8)	(85.0)	(0.2)		(100.0)			
5.0～	55	441	1		497		44	541
(%)	(11.1)	(88.7)	(0.2)		(100.0)			
5.5～	58	433	2		493		40	533
(%)	(11.8)	(87.8)	(0.4)		(100.0)			
6.0～	15	86			101		10	111
(%)	(14.9)	(85.1)			(100.0)			
6.1～	27	218			245		21	266
(%)	(11.0)	(89.0)			(100.0)			
6.5～	43	196	1		240		20	260
(%)	(17.9)	(81.7)	(0.4)		(100.0)			
7.0～	21	134	1		156		16	172
(%)	(13.5)	(85.9)	(0.6)		(100.0)			
7.5～	13	89			102		12	114
(%)	(12.7)	(87.3)			(100.0)			
8.0～	9	59		1	69		8	77
(%)	(13.0)	(85.5)		(1.4)	(100.0)			
8.5～	8	32			40		4	44
(%)	(20.0)	(80.0)			(100.0)			
9.0～	8	41			49		1	50
(%)	(16.3)	(83.7)			(100.0)			
合計	487	3,207	6	2	3,702		376	4,078
(%)	(13.2)	(86.6)	(0.2)	(0.1)	(100.0)			
記載なし	14	43			57		53	110
(%)	(24.6)	(75.4)			(100.0)			
総計	501	3,250	6	2	3,759		429	4,188
(%)	(13.3)	(86.5)	(0.2)	(0.1)	(100.0)			
平均	5.28	5.17	5.77	6.15	5.19		5.01	5.17
標準偏差	1.49	1.46	0.88	2.62	1.46		1.45	1.46

数値下のかっこ内は行方向の合計に対する%です。

##D1629##

(C)Japanese Society for Dialysis Therapy

2016年調査