

表 813 HDF希釈方法 PTH値(pg/mL)別 (オフラインHDF 透析回数週3回) PTH測定法:intact-PTH

PTH値	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
20未満	25	141			166		18	184
(%)	(15.1)	(84.9)			(100.0)			
20～	31	216			247		29	276
(%)	(12.6)	(87.4)			(100.0)			
40～	32	207			239		30	269
(%)	(13.4)	(86.6)			(100.0)			
60～	37	214	2		253		24	277
(%)	(14.6)	(84.6)	(0.8)		(100.0)			
80～	38	257	1		296		26	322
(%)	(12.8)	(86.8)	(0.3)		(100.0)			
100～	38	236			274		24	298
(%)	(13.9)	(86.1)			(100.0)			
120～	32	257	1		290		19	309
(%)	(11.0)	(88.6)	(0.3)		(100.0)			
140～	27	196			223		26	249
(%)	(12.1)	(87.9)			(100.0)			
160～	38	159		1	198		17	215
(%)	(19.2)	(80.3)		(0.5)	(100.0)			
180～	60	377	1		438		45	483
(%)	(13.7)	(86.1)	(0.2)		(100.0)			
240～		4			4			4
(%)		(100.0)			(100.0)			
241～	35	226			261		27	288
(%)	(13.4)	(86.6)			(100.0)			
300～	22	108			130		9	139
(%)	(16.9)	(83.1)			(100.0)			
360～	8	61			69		4	73
(%)	(11.6)	(88.4)			(100.0)			
420～	6	45		1	52		4	56
(%)	(11.5)	(86.5)		(1.9)	(100.0)			
480～	3	12			15			15
(%)	(20.0)	(80.0)			(100.0)			
500～	2	13			15		1	16
(%)	(13.3)	(86.7)			(100.0)			
540～	2	17			19		3	22
(%)	(10.5)	(89.5)			(100.0)			
600～	10	39			49		5	54
(%)	(20.4)	(79.6)			(100.0)			
800～	9	32			41		4	45
(%)	(22.0)	(78.0)			(100.0)			
合計	455	2,817	5	2	3,279		315	3,594
(%)	(13.9)	(85.9)	(0.2)	(0.1)	(100.0)			
記載なし	1	40			41		6	47
(%)	(2.4)	(97.6)			(100.0)			
総計	456	2,857	5	2	3,320		321	3,641
(%)	(13.7)	(86.1)	(0.2)	(0.1)	(100.0)			
平均	182.50	170.72	117.40	296.00	172.35		162.55	171.49
標準偏差	201.65	189.03	67.70	176.78	190.72		157.01	188.01

数値下のかっこ内は行方向の合計に対する%です。

##D1630##

(C)Japanese Society for Dialysis Therapy

2016年調査