

表 819 HDF希釈方法 原疾患別 (オンラインHDF 透析回数週3回)

原疾患	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
慢性糸球体腎炎	16,048	839	1	36	16,924	1	1,061	17,986
(%)	(94.8)	(5.0)	(0.0)	(0.2)	(100.0)			
慢性腎孟腎炎	460	18		1	479		23	502
(%)	(96.0)	(3.8)		(0.2)	(100.0)			
急速進行性糸球体腎炎	353	11		2	366		21	387
(%)	(96.4)	(3.0)		(0.5)	(100.0)			
妊娠腎/妊娠中毒症	328	20		4	352		21	373
(%)	(93.2)	(5.7)		(1.1)	(100.0)			
その他の分類不能の腎炎	236	17		2	255		25	280
(%)	(92.5)	(6.7)		(0.8)	(100.0)			
多発性囊胞腎	1,857	86		2	1,945		121	2,066
(%)	(95.5)	(4.4)		(0.1)	(100.0)			
腎硬化症	4,085	177		14	4,276		258	4,534
(%)	(95.5)	(4.1)		(0.3)	(100.0)			
悪性高血圧	407	14		2	423		17	440
(%)	(96.2)	(3.3)		(0.5)	(100.0)			
糖尿病性腎症	17,053	733		88	17,874		974	18,848
(%)	(95.4)	(4.1)		(0.5)	(100.0)			
SLE腎炎	389	15		5	409		26	435
(%)	(95.1)	(3.7)		(1.2)	(100.0)			
アミロイド腎	64	2			66		7	73
(%)	(97.0)	(3.0)			(100.0)			
痛風腎	165	7			172		13	185
(%)	(95.9)	(4.1)			(100.0)			
先天性代謝異常による腎不全	56	5			61		1	62
(%)	(91.8)	(8.2)			(100.0)			
腎・尿路結核	34	1			35		2	37
(%)	(97.1)	(2.9)			(100.0)			
腎・尿路結石	74	4			78		7	85
(%)	(94.9)	(5.1)			(100.0)			
腎・尿路腫瘍	127	10			137		8	145
(%)	(92.7)	(7.3)			(100.0)			
閉塞性尿路障害	108	2			110		10	120
(%)	(98.2)	(1.8)			(100.0)			
骨髄腫	24	1			25		1	26
(%)	(96.0)	(4.0)			(100.0)			
腎形成不全	116	6			122		13	135
(%)	(95.1)	(4.9)			(100.0)			
不明	4,776	185		13	4,974		238	5,212
(%)	(96.0)	(3.7)		(0.3)	(100.0)			
再導入	482	19		1	502		18	520
(%)	(96.0)	(3.8)		(0.2)	(100.0)			
その他	1,215	55		2	1,272		109	1,381
(%)	(95.5)	(4.3)		(0.2)	(100.0)			
合計	48,457	2,227	1	172	50,857	1	2,974	53,832
(%)	(95.3)	(4.4)	(0.0)	(0.3)	(100.0)			
記載なし								
(%)								
総計	48,457	2,227	1	172	50,857	1	2,974	53,832
(%)	(95.3)	(4.4)	(0.0)	(0.3)	(100.0)			

数値下の括弧内は行方向の合計に対する%です。

##D1713##

(C)Japanese Society for Dialysis Therapy

2016年調査