

表 827 HDF希釈方法 single pool Kt/V(Kt/Vsp)別（オンラインHDF 男性 透析回数週3回 透析歴2年以上）

single pool Kt/V (Kt/Vsp)	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
0.4未満	16	2			18		2	20
(%)	(88.9)	(11.1)			(100.0)			
0.4～	13	3			16		1	17
(%)	(81.3)	(18.8)			(100.0)			
0.6～	63	3			66		2	68
(%)	(95.5)	(4.5)			(100.0)			
0.8～	558	32		2	592		42	634
(%)	(94.3)	(5.4)		(0.3)	(100.0)			
1.0～	2,844	119		6	2,969		170	3,139
(%)	(95.8)	(4.0)		(0.2)	(100.0)			
1.2～	7,330	352		22	7,704		389	8,093
(%)	(95.1)	(4.6)		(0.3)	(100.0)			
1.4～	7,489	355		12	7,856		439	8,295
(%)	(95.3)	(4.5)		(0.2)	(100.0)			
1.6～	3,690	173		4	3,867		233	4,100
(%)	(95.4)	(4.5)		(0.1)	(100.0)			
1.8～	1,472	96		1	1,569		93	1,662
(%)	(93.8)	(6.1)		(0.1)	(100.0)			
2.0～	793	45			838		52	890
(%)	(94.6)	(5.4)			(100.0)			
合計	24,268	1,180		47	25,495		1,423	26,918
(%)	(95.2)	(4.6)		(0.2)	(100.0)			
記載なし	2,578	86	1	21	2,686		215	2,901
(%)	(96.0)	(3.2)	(0.0)	(0.8)	(100.0)			
総計	26,846	1,266	1	68	28,181		1,638	29,819
(%)	(95.3)	(4.5)	(0.0)	(0.2)	(100.0)			
平均	1.45	1.46		1.36	1.45		1.46	1.45
標準偏差	0.26	0.28		0.19	0.27		0.27	0.27

数値下のかっこ内は行方向の合計に対する%です。

##D1723##

(C)Japanese Society for Dialysis Therapy

2016年調査