

表 836 HDF希釈方法 透析前リン濃度(mg/dL)別 (オンラインHDF 透析回数週3回)

透析前リン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3未満	1,416	49		10	1,475		99	1,574
(%)	(96.0)	(3.3)		(0.7)	(100.0)			
3.0~	1,865	85		13	1,963	1	94	2,058
(%)	(95.0)	(4.3)		(0.7)	(100.0)			
3.5~	3,538	168		10	3,716		175	3,891
(%)	(95.2)	(4.5)		(0.3)	(100.0)			
4.0~	5,511	245		12	5,768		320	6,088
(%)	(95.5)	(4.2)		(0.2)	(100.0)			
4.5~	7,133	303		16	7,452		404	7,856
(%)	(95.7)	(4.1)		(0.2)	(100.0)			
5.0~	7,324	350		17	7,691		420	8,111
(%)	(95.2)	(4.6)		(0.2)	(100.0)			
5.5~	6,569	346		17	6,932		385	7,317
(%)	(94.8)	(5.0)		(0.2)	(100.0)			
6.0~	1,119	53		4	1,176		69	1,245
(%)	(95.2)	(4.5)		(0.3)	(100.0)			
6.1~	3,825	189	1	10	4,025		238	4,263
(%)	(95.0)	(4.7)	(0.0)	(0.2)	(100.0)			
6.5~	3,497	153		7	3,657		213	3,870
(%)	(95.6)	(4.2)		(0.2)	(100.0)			
7.0~	2,182	94		10	2,286		134	2,420
(%)	(95.5)	(4.1)		(0.4)	(100.0)			
7.5~	1,310	67		2	1,379		88	1,467
(%)	(95.0)	(4.9)		(0.1)	(100.0)			
8.0~	816	45		2	863		51	914
(%)	(94.6)	(5.2)		(0.2)	(100.0)			
8.5~	518	32		1	551		33	584
(%)	(94.0)	(5.8)		(0.2)	(100.0)			
9.0~	784	34		1	819		45	864
(%)	(95.7)	(4.2)		(0.1)	(100.0)			
合計	47,407	2,213	1	132	49,753	1	2,768	52,522
(%)	(95.3)	(4.4)	(0.0)	(0.3)	(100.0)			
記載なし	1,050	14		40	1,104		206	1,310
(%)	(95.1)	(1.3)		(3.6)	(100.0)			
総計	48,457	2,227	1	172	50,857	1	2,974	53,832
平均	5.35	5.41	6.10	5.07	5.35	3.00	5.39	5.35
標準偏差	1.42	1.39		1.57	1.42		1.46	1.42

数値下の括弧内は行方向の合計に対する%です。

##D1730##

(C)Japanese Society for Dialysis Therapy

2016年調査