

表 843 HDF希釈方法 原疾患別 (IHDF 透析回数週3回)

原疾患	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
慢性糸球体腎炎	222	53	47	1,013	1,335	4	1,320	2,659
(%)	(16.6)	(4.0)	(3.5)	(75.9)	(100.0)			
慢性腎盂腎炎	10	1	1	26	38		37	75
(%)	(26.3)	(2.6)	(2.6)	(68.4)	(100.0)			
急速進行性糸球体腎炎	6	1		26	33		30	63
(%)	(18.2)	(3.0)		(78.8)	(100.0)			
妊娠腎/妊娠中毒症	2	1	1	18	22		16	38
(%)	(9.1)	(4.5)	(4.5)	(81.8)	(100.0)			
その他の分類不能の腎炎	4	2	1	34	41		11	52
(%)	(9.8)	(4.9)	(2.4)	(82.9)	(100.0)			
多発性嚢胞腎	37	6		159	202	1	154	357
(%)	(18.3)	(3.0)		(78.7)	(100.0)			
腎硬化症	118	22		396	536		540	1,076
(%)	(22.0)	(4.1)		(73.9)	(100.0)			
悪性高血圧	4			48	52		39	91
(%)	(7.7)			(92.3)	(100.0)			
糖尿病性腎症	380	71	27	1,682	2,160	4	2,042	4,206
(%)	(17.6)	(3.3)	(1.3)	(77.9)	(100.0)			
SLE腎炎	5	2	1	13	21		30	51
(%)	(23.8)	(9.5)	(4.8)	(61.9)	(100.0)			
アミロイド腎				3	3		11	14
(%)				(100.0)	(100.0)			
痛風腎	1			10	11		14	25
(%)	(9.1)			(90.9)	(100.0)			
先天性代謝異常による腎不全	2			2	4		3	7
(%)	(50.0)			(50.0)	(100.0)			
腎・尿路結核				4	4		3	7
(%)				(100.0)	(100.0)			
腎・尿路結石	1			8	9		11	20
(%)	(11.1)			(88.9)	(100.0)			
腎・尿路腫瘍	1			10	11		18	29
(%)	(9.1)			(90.9)	(100.0)			
閉塞性尿路障害	2	1		9	12		11	23
(%)	(16.7)	(8.3)		(75.0)	(100.0)			
骨髄腫				1	1		1	2
(%)				(100.0)	(100.0)			
腎形成不全	2			9	11		9	20
(%)	(18.2)			(81.8)	(100.0)			
不明	81	6	1	463	551		389	940
(%)	(14.7)	(1.1)	(0.2)	(84.0)	(100.0)			
再導入	2		3	37	42		19	61
(%)	(4.8)		(7.1)	(88.1)	(100.0)			
その他	17	3		103	123		118	241
(%)	(13.8)	(2.4)		(83.7)	(100.0)			
合計	897	169	82	4,074	5,222	9	4,826	10,057
(%)	(17.2)	(3.2)	(1.6)	(78.0)	(100.0)			
記載なし								
(%)								
総計	897	169	82	4,074	5,222	9	4,826	10,057
(%)	(17.2)	(3.2)	(1.6)	(78.0)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

##D1813##

(C)Japanese Society for Dialysis Therapy

2016年調査