

表 861 HDF希釈方法 透析前リン濃度(mg/dL)別 (IHDF 透析回数週3回)

透析前リン濃度	前希釈	後希釈	前・後希釈	その他	合計	不明	記載なし	総計
3未満	36	5	1	108	150		223	373
(%)	(24.0)	(3.3)	(0.7)	(72.0)	(100.0)			
3.0~	36	6	1	187	230	1	269	500
(%)	(15.7)	(2.6)	(0.4)	(81.3)	(100.0)			
3.5~	89	4	4	340	437	1	417	855
(%)	(20.4)	(0.9)	(0.9)	(77.8)	(100.0)			
4.0~	110	17	10	481	618	1	652	1,271
(%)	(17.8)	(2.8)	(1.6)	(77.8)	(100.0)			
4.5~	129	23	14	562	728	1	724	1,453
(%)	(17.7)	(3.2)	(1.9)	(77.2)	(100.0)			
5.0~	144	22	11	585	762		711	1,473
(%)	(18.9)	(2.9)	(1.4)	(76.8)	(100.0)			
5.5~	117	29	11	550	707		611	1,318
(%)	(16.5)	(4.1)	(1.6)	(77.8)	(100.0)			
6.0~	22	3	3	89	117		83	200
(%)	(18.8)	(2.6)	(2.6)	(76.1)	(100.0)			
6.1~	54	17	3	338	412	3	309	724
(%)	(13.1)	(4.1)	(0.7)	(82.0)	(100.0)			
6.5~	60	19	8	273	360	1	269	630
(%)	(16.7)	(5.3)	(2.2)	(75.8)	(100.0)			
7.0~	31	4	8	179	222	1	198	421
(%)	(14.0)	(1.8)	(3.6)	(80.6)	(100.0)			
7.5~	30	8	5	114	157		87	244
(%)	(19.1)	(5.1)	(3.2)	(72.6)	(100.0)			
8.0~	20	2		70	92		62	154
(%)	(21.7)	(2.2)		(76.1)	(100.0)			
8.5~	7	2	1	39	49		24	73
(%)	(14.3)	(4.1)	(2.0)	(79.6)	(100.0)			
9.0~	6	3	2	65	76		68	144
(%)	(7.9)	(3.9)	(2.6)	(85.5)	(100.0)			
合計	891	164	82	3,980	5,117	9	4,707	9,833
(%)	(17.4)	(3.2)	(1.6)	(77.8)	(100.0)			
記載なし	6	5		94	105		119	224
(%)	(5.7)	(4.8)		(89.5)	(100.0)			
総計	897	169	82	4,074	5,222	9	4,826	10,057
平均	5.18	5.56	5.62	5.32	5.31	5.43	5.10	5.21
標準偏差	1.38	1.44	1.37	1.42	1.42	1.44	1.42	1.42

数値下の括弧内は行方向の合計に対する%です。

##D1829##

(C)Japanese Society for Dialysis Therapy

2016年調査